

#12713 Store at -20C

uPAR (D7X2N) Rabbit mAb**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IF-IC	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 35-65	Source/Isotype: Rabbit IgG	UniProt ID: #Q03405	Entrez-Gene Id: 5329
----------------------------	------------------	----------------------------	--------------------	-------------------------------	------------------------	-------------------------

Product Usage Information**Application**Western Blotting
Immunofluorescence (Immunocytochemistry)**Dilution**1:1000
1:50**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

For a carrier free (BSA and azide free) version of this product see product #45547.

Specificity / Sensitivity

uPAR (D7X2N) Rabbit mAb recognizes endogenous levels of total uPAR protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Arg303 of human uPAR protein.

Background

The human urokinase-type plasminogen activator receptor (uPAR) is a 55-65 kDa, highly glycosylated, GPI-anchored cell surface receptor (the deglycosylated protein is 35 kDa) (1-3). It is a central player in the plasminogen activation pathway. uPAR binds with high affinity to a serine protease urokinase-type plasminogen activator (uPA) and converts plasminogen to its active form plasmin in a spatially restricted manner on the cell surface (4). Plasmin further carries out the activation of uPA, which is inhibited by serpins, such as plasminogen activator inhibitors (5). Therefore, uPAR plays a key role in regulating extracellular proteolysis. In addition, uPAR plays an important role in regulating cell proliferation, adhesion and mobility (6,7). Research studies have shown that overexpression of uPAR is found in various cancer cells and tissues (8,9).

Background References

1. Nielsen, L.S. et al. (1988) *J Biol Chem* 263, 2358-63.
2. Behrendt, N. et al. (1990) *J Biol Chem* 265, 6453-60.
3. Roldan, A.L. et al. (1990) *EMBO J* 9, 467-74.
4. Ellis, V. et al. (1991) *J Biol Chem* 266, 12752-8.
5. Ellis, V. et al. (1990) *J Biol Chem* 265, 9904-8.
6. Liu, D. et al. (2002) *Cancer Cell* 1, 445-57.
7. Waltz, D.A. et al. (1997) *J Clin Invest* 100, 58-67.
8. Blasi, F. and Sidenius, N. (2010) *FEBS Lett* 584, 1923-30.
9. Mazar, A.P. et al. (2011) *Curr Pharm Des* 17, 1970-8.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key**WB:** Western Blotting **IF-IC:** Immunofluorescence (Immunocytochemistry)**Cross-Reactivity Key**

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
 All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.