mAb

2936 Store at -20C

Orders:	877-616-CELL (2355) orders@cellsignal.com			
Support:	877-678-TECH (8324)			
Web:	info@cellsignal.com cellsignal.com			

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

Galectin-1/LGALS1 (D608T) Rabbit

Applications: WB, IF-IC	Reactivity: H M R Mk	Sensitivity: Endogenous	<b>MW (kDa):</b> 15	Source/Isotype: Rabbit IgG	UniProt ID: #P09382	Entrez-Gene Id: 3956		
Product Usage Information	We	plication estern Blotting munofluorescence (In	nmunocytochen	nistry)		<b>Dilution</b> 1:1000 1:50		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.						
Specificity / Sensit		Galectin-1/LGALS1 (D6O8T) Rabbit mAb recognizes endogenous levels of total galectin-1/LGALS1 protein.						
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human galectin-1/LGALS1 protein.						
Background	ace gale regi has exp dev func sho	Galectins are a family of β-galactose binding proteins that are characterized by their affinity for poly-N- acetyllactosamine-enriched glycoconjugates and their carbohydrate-binding site (1,2). Members of the galectin family have been implicated in a variety of biological functions including cell adhesion (3), growth regulation (4), cytokine production (5), T cell apoptosis (6), and immune responses (7). Galectin-1/LGALS1 has been shown to be expressed in a wide range of tissues and cell types. The level and pattern of expression of galectin-1 have been shown to change during development (8). In addition to a role in developmental processes, galectin-1 has been shown to be involved in central immune tolerance and may function in tumorigenesis by modulating the immune response to the tumor (9,10). Research studies have shown that galectin-1 expression is increased in several human cancers, suggesting a correlation with metastatic potential (10).						
Background Refer	2. B 3. C 4. V 5. F 6. P 7. C 8. P 9. v	arondes, S.H. et al. (: arondes, S.H. et al. (: offner, H. et al. (1990) Vells, V. and Mallucci, iler, A. et al. (2009) A. erillo, N.L. et al. (199 cooper, D.N. et al. (199 uche, A.C. et al. (199 an den Brûle, F. et al. alatino, M. et al. (200	1994) J Biol Chu J Neuroimmund L. (1991) Cell 6 rthritis Rheum 6 5) Nature 378, 7 91) J Cell Biol 1 6) Dev Biol 179 (2003) Lab Inve	em 269, 20807-10. ol 28, 177-84. 54, 91-7. 50, 1604-14. 736-9. .15, 1437-48. 1, 274-87. est 83, 377-86.				
Species Reactivity	Spec	Species reactivity is determined by testing in at least one approved application (e.g., western blot).						
Western Blot Buffe		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications Key	WB	WB: Western Blotting IF-IC: Immunofluorescence (Immunocytochemistry)						
Cross-Reactivity K	X: X	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected						
Trademarks and Patents	All o	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.						
Limited Uses								

Galectin-1/LGALS1 (D608T) Rabbit mAb (#12936) Datasheet Without Images Cell Signaling Technology

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.