

#13381 Store at -20C

Artemis (D7O8V) Rabbit mAb

Cell Signaling
TECHNOLOGY®Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H Mk	Endogenous	90	Rabbit IgG	#Q96SD1	64421

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Immunoprecipitation

1:200

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Artemis (D7O8V) Rabbit mAb recognizes endogenous levels of total artemis protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro367 of human artemis protein.

Background

DNA double-strand breaks (DSBs) are potentially hazardous lesions that can be induced by ionizing radiation (IR), radiomimetic chemicals, or DNA replication inhibitors. Cells recognize and repair DSBs via two distinct but partly overlapping signaling pathways, nonhomologous end joining (NHEJ) and homologous recombination (HR). DNA repair via the HR pathway is restricted to S and G2 phases of the cell cycle, while NHEJ can occur during any phase. Defects in both pathways have been associated with human disease, including cancer (1).

Artemis is a ubiquitously expressed NHEJ factor that exhibits endonuclease activity. Artemis functions in DNA repair by promoting nonhomologous end joining (2), as well as in cell cycle checkpoint control through ATM/ATR signaling (3).

NHEJ machinery is also utilized in V(D)J recombination, a process that generates diversity in immunoglobulin and T cell receptor genes, and artemis is a key factor in this process (4,5). Mutations in the corresponding artemis gene (*DCLRE1C*) are associated with a radiosensitive type of severe combined immunodeficiency (SCID) in humans (6,7).

Background References

- Hartlerode, A.J. and Scully, R. (2009) *Biochem J* 423, 157-68.
- Kurosawa, A. et al. (2013) *PLoS One* 8, e72253.
- Zhang, X. et al. (2004) *Mol Cell Biol* 24, 9207-20.
- Mansilla-Soto, J. and Cortes, P. (2003) *J Exp Med* 197, 543-7.
- Ma, Y. et al. (2002) *Cell* 108, 781-94.
- Noordzij, J.G. et al. (2003) *Blood* 101, 1446-52.
- Li, L. et al. (2002) *J Immunol* 168, 6323-9.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster
X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse
GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.