

#13571 Store at -20°C

Phospho-WIP12 (Ser413) Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R	Endogenous	49	Rabbit	#Q9Y4P8	26100

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Phospho-WIP12 (Ser413) recognizes endogenous levels of WIP12 protein only when phosphorylated at Ser413.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser413 of human WIP12 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Autophagy is a catabolic process for the autophagosomic-lysosomal degradation of bulk cytoplasmic contents (1,2). It is generally activated by conditions of nutrient deprivation but is also associated with a number of physiological processes including development, differentiation, neurodegeneration, infection, and cancer (3). The molecular machinery of autophagy was largely discovered in yeast and is directed by a number of autophagy-related (Atg) genes.

Vacuolar trafficking and autophagy are controlled by the class III type phosphoinositide 3-kinase (PI3K) Vps34, which generates phosphoinositide-3-phosphate (PtdIns3P) (4,5). Atg18 and Atg21 are two related WD-repeat proteins that bind PtdIns3P via a conserved Phe-Arg-Arg-Gly motif (6,7). It has been shown that Atg18 binds to Atg2 and that this complex is directed to vacuolar membranes by its interaction with PtdIns3P (8). Human orthologs of Atg18 and Atg21 were identified as members of the WD-repeat protein Interacting with Phosphoinositides (WIP1) family (9-11). WIP1 (also called WIP149) and WIP12 have been shown to translocate from several vacuolar compartments to LC3-positive autophagosomes during autophagy; this translocation may be used as an autophagy marker (10).~

Phospho-WIP12 (Ser413) Antibody is directed at a site that was identified at Cell Signaling Technology (CST) using PhosphoScan, CST's LC-MS/MS platform for modification site discovery. Phosphorylation at Ser413 was discovered using Phospho-MAPK/CDK Substrate motif antibody. Please visit PhosphoSitePlus, CST's modification site knowledgebase, at www.phosphosite.org for more information.

Background References

- Reggiori, F. and Klionsky, D.J. (2002) *Eukaryot Cell* 1, 11-21.
- Codogno, P. and Meijer, A.J. (2005) *Cell Death Differ* 12 Suppl 2, 1509-18.
- Levine, B. and Yuan, J. (2005) *J Clin Invest* 115, 2679-88.
- Corvera, S. (2001) *Traffic* 2, 859-66.
- Yan, Y. and Backer, J.M. (2007) *Biochem Soc Trans* 35, 239-41.
- Krick, R. et al. (2006) *FEBS Lett* 580, 4632-8.
- Strømhaug, P.E. et al. (2004) *Mol Biol Cell* 15, 3553-66.
- Obara, K. et al. (2008) *J Biol Chem* 283, 23972-80.
- Jeffries, T.R. et al. (2004) *Mol Biol Cell* 15, 2652-63.
- Proikas-Cezanne, T. et al. (2007) *FEBS Lett* 581, 3396-404.
- Polson, H.E. et al. (2010) *Autophagy* 6, 506-22.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
PhosphoScan is a registered trademark of Cell Signaling Technology, Inc.
PhosphoSitePlus is a registered trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.