

#13805 Store at -20C

XAF1 (E1E4O) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H	Endogenous	32, 38	Rabbit IgG	#Q6GPH4	54739

Product Usage Information

Application

Western Blotting

Immunoprecipitation

Dilution

1:1000

1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

XAF1 (E1E4O) Rabbit mAb recognizes endogenous levels of total XAF1 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Val210 of human XAF1 protein.

Background

X-linked inhibitor of apoptosis (XIAP)-associated factor 1 (XAF1) is a zinc finger protein that antagonizes the anti-apoptotic activity of XIAP (1,2). XIAP, a member of the inhibitor of apoptosis (IAP) family, inhibits apoptosis by direct inhibition of caspases (3; reviewed in 4). XAF1 is widely expressed in normal tissues, with highest levels in the heart and ovary, but is mostly reduced in cancer lines (1,2). Expression of XAF1 can be induced by interferons via Stat transcriptional activity (5-7). The levels of XAF1 have been shown to be inversely correlated with p53, and p53 is directly responsible for inhibiting XAF1 transcription (8,9). A number of studies have shown that XAF1 can function as a tumor suppressor protein, and decreased levels of XAF1 are found in a variety of different cancers (10-13). Research studies suggest that expression of XAF1 may be a prognostic biomarker for some cancers (14-16).

Background References

1. Fong, W.G. et al. (2000) *Genomics* 70, 113-22.
2. Liston, P. et al. (2001) *Nat Cell Biol* 3, 128-33.
3. Deveraux, Q.L. et al. (1997) *Nature* 388, 300-4.
4. Goyal, L. (2001) *Cell* 104, 805-8.
5. Leaman, D.W. et al. (2002) *J Biol Chem* 277, 28504-11.
6. Sun, Y. et al. (2008) *Cancer Lett* 260, 62-71.
7. Bai, Y. et al. (2008) *J Biol Chem* 283, 6832-42.
8. Zhang, W. et al. (2010) *Int J Oncol* 36, 1031-7.
9. Zou, B. et al. (2012) *Mol Carcinog* 51, 422-32.
10. Byun, D.S. et al. (2003) *Cancer Res* 63, 7068-75.
11. Ng, K.C. et al. (2004) *J Invest Dermatol* 123, 1127-34.
12. Ma, T.L. et al. (2005) *Chin J Dig Dis* 6, 10-4.
13. Zou, B. et al. (2006) *Gastroenterology* 131, 1835-43.
14. Huang, J. et al. (2010) *Cancer Sci* 101, 559-67.
15. Chen, Y.B. et al. (2011) *Chin Med J (Engl)* 124, 3238-43.
16. Wang, Y. et al. (2012) *Regul Pept* 178, 36-42.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.