

#14181 Store at -20°C

## CLPP Antibody

Cell Signaling  
TECHNOLOGY®Orders: 877-616-CELL (2355)  
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H	Endogenous	28	Rabbit	#Q16740	8192

## Product Usage Information

## Application

Western Blotting

## Dilution

1:1000

Immunoprecipitation

1:50

## Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. *Do not aliquot the antibody.*

## Specificity / Sensitivity

CLPP Antibody recognizes endogenous levels of total CLPP protein.

## Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro263 of human CLPP protein. Antibodies are purified by protein A and peptide affinity chromatography.

## Background

The proteolytic component, Tetradecameric Peptidase (CLpP), is a hexamer in one of four ATP-dependent mitochondrial proteases (CLpXP). CLPP, one of the proteases, is an endopeptidase that is highly conserved among prokaryotes and eukaryotes, both at the level of amino acid sequence and quaternary structure. The active unit of CLPP is a barrel-shaped tetradecamer. Proteolysis of larger substrates is initiated by caseinolytic peptidase X (CLPX) which unfolds specific protein substrates. The unfolded polypeptide chain translocates into the CLPP proteolytic chamber for protein degradation within the interior chamber of mitochondria (1). Recessive mutations in *CLPP* cause Perrault Syndrome, a heterogeneous condition characterized by sensorineural hearing loss and ovarian failure (2).

Mutations in Parkin or PINK1 cause recessively inherited Parkinson's disease. In healthy mitochondria, PINK1 is rapidly degraded by mitochondrial proteases and the proteasome. Upon mitochondrial depolarization, PINK1 accumulates on the mitochondrial surface, recruits Parkin from the cytosol, and initiates mitophagy. The mitochondrial proteases MPP, PARL, m-AAA and CLPP have been implicated in PINK1 degradation and cleavage (3).

## Background References

1. Yu, A.Y. and Houry, W.A. (2007) *FEBS Lett* 581, 3749-57.
2. Jenkinson, E.M. et al. (2013) *Am J Hum Genet* 92, 605-13.
3. Greene, A.W. et al. (2012) *EMBO Rep* 13, 378-85.

## Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

## Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

## Applications Key

**WB:** Western Blotting **IP:** Immunoprecipitation

## Cross-Reactivity Key

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

## Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
 All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.