$522\,$ Store at -200

ARFGAP1 (D9A4V) Rabbit mAb



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support:

877-678-TECH (8324)

Web:

info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 45	Source/Isotype: Rabbit IgG	UniProt ID: #Q8N6T3	Entrez-Gene Id: 55738
Product Usage Information	Ар	plication		Dilution		
	We	estern Blotting		1:1000		
	Im	munoprecipitation			1:50	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20° C. Do not aliquot the antibody.				
Specificity / Sensiti	vity ARI	ARFGAP1 (D9A4V) Rabbit mAb recognizes endogenous levels of total ARFGAP1 protein.				
Source / Purificatio		noclonal antibody is dues surrounding Th		nunizing animals with a ARFGAP1 protein.	synthetic peptide corre	esponding to
Background	ves forn prot acti stuc prot GTI	ADP-ribosylation factor GTPase activating protein 1 (ARFGAP1) is a Golgi-localized protein that regulates vesicle formation and membrane trafficking (1). ARFGAP1 initiates cargo selection and COP1 vesicle formation by stimulating GTP hydrolysis of ADP-ribosylation factor ARF1 (2). This GTPase activating protein initiates vesicle transport by coupling vesicle formation with cargo sorting (3). ARFGAP1 plays an active role in the Golgi-to-ER retrograde, intra-Golgi, and trans-Golgi trafficking networks (1). Research studies indicate that ARFGAP1 can act as a GTPase activating protein for LRRK2, a large multifunction protein whose genetic mutations are associated with Parkinson's disease (4). ARFGAP1 regulates GTPase activity and promotes the kinase activity of LRRK2, which suggests some potential as a promising target for study of LRRK2 mediated neurodegeneration (4).				
Background Refere	2. K 3. Y	noue, H. and Randa: Lartberg, F. et al. (20 Kang, J.S. et al. (2000 Stafa, K. et al. (2012)	10) J Biol Chem 2) J Cell Biol 159	285, 36709-20. , 69-78.		

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry

milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner ARFGAP1 (D9A4V) Rabbit mAb (#14522) Datasheet Without Images Cell Signaling Technology

that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.