

#14674 Store at -20°C

STAM2 Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H Mk	Endogenous	70	Rabbit	#O75886	10254

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

STAM2 Antibody recognizes endogenous levels of total STAM2 protein. This antibody does not cross-react with STAM1.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human STAM2 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Signal transducing adaptor molecule 2 (STAM2) is a ubiquitously expressed STAM family adaptor protein and an integral component of the ESCRT-0 complex. Similar to STAM1, STAM2 possesses a single SH3 domain and an immunoreceptor tyrosine-based activation motif (ITAM). Following activation of multiple growth factor and cytokine cell surface receptors, the STAM2 protein undergoes tyrosine phosphorylation and potentiates mitogenic signals driven by these receptors (1,2). Research studies demonstrate that STAM2 is localized to complexes containing Eps15, Hrs, and STAM1 proteins on early endosome membranes. A tandem, amino-terminal VHS (Vps27/Hrs/STAM) domain and UIM (ubiquitin-interacting) motif within STAM2 facilitate STAM2 interaction with ubiquitinated cargo proteins, suggesting that this adaptor participates in the endosomal sorting of ubiquitinated proteins targeted for lysosomal degradation (3-6). Gene targeting studies have revealed an indispensable role for STAM2 in T-cell development (7).

Background References

- Endo, K. et al. (2000) *FEBS Lett* 477, 55-61.
- Pandey, A. et al. (2000) *J Biol Chem* 275, 38633-9.
- Mizuno, E. et al. (2003) *Mol Biol Cell* 14, 3675-89.
- Bache, K.G. et al. (2003) *J Biol Chem* 278, 12513-21.
- Takata, H. et al. (2000) *Genes Cells* 5, 57-69.
- Mizuno, E. et al. (2004) *J Biochem* 135, 385-96.
- Yamada, M. et al. (2002) *Mol Cell Biol* 22, 8648-58.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.