

#14696 Store at -20C

CBX8 (D2O8C) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP, ChIP, ChIP-seq, C&R, C&T	H M R Mk	Endogenous	43, 48	Rabbit IgG	#Q9HC52	57332

Product Usage Information

For optimal ChIP and ChIP-seq results, use 10 µl of antibody and 10 µg of chromatin (approximately 4 x 10⁶ cells) per IP. This antibody has been validated using SimpleChIP® Enzymatic Chromatin IP Kits.

The CUT&RUN dilution was determined using CUT&RUN Assay Kit #86652.

The CUT&Tag dilution was determined using CUT&Tag Assay Kit #77552.

Application	Dilution
Western Blotting	1:1000
Immunoprecipitation	1:200
Chromatin IP	1:50
Chromatin IP-seq	1:50
CUT&RUN	1:50
CUT&Tag	1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

CBX8 (D2O8C) Rabbit mAb recognizes endogenous levels of total CBX8 protein.

Species predicted to react based on 100% sequence homology:

Hamster, Bovine

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly344 of human CBX8 protein.

Background

The polycomb group (PcG) proteins contribute to the maintenance of cell identity, stem cell self-renewal, cell cycle regulation and oncogenesis by maintaining the silenced state of genes that promote cell lineage specification, cell death and cell-cycle arrest (1-4). PcG proteins exist in two complexes that cooperate to maintain long-term gene silencing through epigenetic chromatin modifications: PRC1 and PRC2. PRC1 is a multi-subunit protein complex consisting of a combination of five core protein families: CBX, RING1, PHC, PCGF, and RYPB (5-7). Different combinations of protein family members lead to a diverse array of PRC1 complexes with distinct functions (8). At least two distinct classes of PRC1 complexes have been defined. The first class, known as canonical PRC1, contains RING1, PHC, PCGF and CBX protein subunits, but not RYPB (5-8). This class of PRC1 complexes requires PRC2 and H3K27Me3 for proper recruitment to target genes. CBX proteins mediate recruitment by binding to H3K27Me3. CBX8 in particular is required for repression of many lineage-specific genes during differentiation of hematopoietic stem cells and may play a role in activation of lineage-specific genes during differentiation of embryonic stem cells (9,10). The second class, known as variant PRC1, contains RYPB instead of CBX proteins (5-8). RYPB-containing PRC1 is recruited to chromatin independently of PRC2 and H3K27Me3. These variant PRC1 complexes can function independently of PRC2, or in some cases function upstream to recruit PRC2 complex to target genes.

Background References

1. Boyer, L.A. et al. (2006) *Nature* 441, 349-53.
2. Lee, T.I. et al. (2006) *Cell* 125, 301-13.
3. Park, I.K. et al. (2003) *Nature* 423, 302-5.
4. Molofsky, A.V. et al. (2003) *Nature* 425, 962-7.
5. Tavares, L. et al. (2012) *Cell* 148, 664-78.

6. Gao, Z. et al. (2012) *Mol Cell* 45, 344-56.
7. Blackledge, N.P. et al. (2014) *Cell* 157, 1445-59.
8. Luis, N.M. et al. (2012) *Cell Stem Cell* 11, 16-21.
9. Klauke, K. et al. (2013) *Nat Cell Biol* 15, 353-62.
10. Creppe, C. et al. (2014) *PLoS Genet* 10, e1004851.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation **ChIP:** Chromatin IP **ChIP-seq:** Chromatin IP-seq
C&R: CUT&RUN **C&T:** CUT&Tag

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
SimpleChIP is a registered trademark of Cell Signaling Technology, Inc.
XP is a registered trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.