74775 Store at -20C

Pontin/RUVBL1 (D1L8J) XP® Rabbit mAb Orders: 877-616-CELL (2355) orders@cellsignal.com

> Web: info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

	Reactivity: Sensitivit H M R Mk Endogeno		Source/Isotype: Rabbit IgG	UniProt ID: #Q9Y265	Entrez-Gene Id 8607	
Product Usage Information	For optimal ChIP r This antibody has	For optimal ChIP results, use 10 μ l of antibody and 10 μ g of chromatin (approximately 4 x 10 ⁶ cells) per IP. This antibody has been validated using SimpleChIP [®] Enzymatic Chromatin IP Kits.				
	Application			Dilution		
	Western Blotting			1:1000		
	Chromatin IP	Chromatin IP 1:50				
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less t 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity / Sensitivi	ity Pontin/RUVBL1 (D	1L8J) XP [®] Rabbit m	Ab recognizes endogenc	ous levels of total Pon	tin/RUVBL1 protein.	
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Glu428 of human Pontin/RUVBL1 protein.			esponding to	
Background	diverse cellular ac proteins that drive essential compone SRCAP, and Uri1, transcriptional co-r	ivities) superfamily of branch migration of H ints of chromatin rem which play key roles i	closely related members f proteins, and are putati folliday junctions (1). Re odeling and modification in regulating gene transo I Pontin have both been hin, and E1A (2-7).	vely homologous to be ptin and Pontin function complexes, such as cription (1,2). In their c	acterial RuvB on together as INO80, TIP60, apacity as essential	
Background Referen	2. Gallant, P. (2007 3. Huber, O. et al. 4. Kim, J.H. et al. (5. Bauer, A. et al. (6. Wood, M.A. et al.	a, A. (2009) Mol Cell 7) Trends Cell Biol 17 (2008) Cancer Res 68 2005) Nature 434, 92 2000) EMBO J 19, 61 I. (2000) Mol Cell 5, 3 II. (2002) Oncogene 2	, 187-92. 8, 6873-6. 11-6. 121-30. 321-30.			
Species Reactivity	Species reactivity is	Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key	WB: Western Blott	ing ChIP: Chromatin	IP			
Cross-Reactivity Key	X: Xenopus Z: zeb	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected				
Trademarks and Patents	SimpleChIP is a reg XP is a registered t	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. SimpleChIP is a registered trademark of Cell Signaling Technology, Inc. XP is a registered trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.				
Limited Uses	following terms app	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in				

Pontin/RUVBL1 (D1L8J) XP® Rabbit mAb (#74775) Datasheet Without Images Cell Signaling Technology

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com For Research Use Only. Not for Use in Diagnostic Procedures.