499 Store at -200

POMC (D3R1U) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Reactivity:	Sensitivity: Endogenous	MW (kDa): 10, 12, 28, 32	Source/Isotype: Rabbit IgG	UniProt ID: #P01189	Entrez-Gene Id: 5443
Application					Dilution
Wes	Western Blotting				1:1000
Imm	nunoprecipitation				1:50
Imm	nunofluorescence ((Frozen)			1:200
Imm	Immunofluorescence (Immunocytochemistry)				1:800
		\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	,,	10 , 0)	cerol and less than
vity POM	C (D3R1U) Rabbi	t mAb recognizes	endogenous levels of to	otal POMC protein.	
o Hum % y:	an, Rat, Bovine				
	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro69 of human POMC protein.				
proce MSH regul	Pro-opiomelanocortin (POMC) is a precursor protein expressed in the pituitary and the brain where it is processed into several peptide hormones and neuropeptides. Among these peptides are ACTH, α - and β -MSH, β -and γ -LPH, CLIP, β -endorphin, and N-POMC (1). POMC is involved in hypothalamic circuits regulating feeding behavior and POMC-producing neurons promote satiety (2). POMC neurons are also the target of leptin and insulin for the promotion of the browning of white fat (3).				
2. Ya	 Chretien, M. and Mbikay, M. (2016) J Mol Endocrinol, T49-62. doi: 10.1530/JME-15-0261. Yang, Y. et al. (2011) Cell 146, 992-1003. Dodd, G.T. et al. (2015) Cell 160, 88-104. 				
	App Wes Imm Imm Supp 0.029 vity POM Hum % y: n Monoresid Pro-co proce MSH regul the ta ences 1. Ch 2. Ya	Application Western Blotting Immunoprecipitation Immunofluorescence (Immunofluorescence (Supplied in 10 mM sod 0.02% sodium azide. S POMC (D3R1U) Rabbi Human, Rat, Bovine Monoclonal antibody is residues surrounding P Pro-opiomelanocortin (processed into several MSH, β-and γ-LPH, CL regulating feeding behat the target of leptin and 1. Chretien, M. and Mb 2. Yang, Y. et al. (2011)	Application Western Blotting Immunoprecipitation Immunofluorescence (Frozen) Immunofluorescence (Immunocytochem Supplied in 10 mM sodium HEPES (pH 7 0.02% sodium azide. Store at –20°C. Do POMC (D3R1U) Rabbit mAb recognizes of the target of leptin and insulin for the prorection. Monoclonal antibody is produced by immunosidues surrounding Pro69 of human PC Pro-opiomelanocortin (POMC) is a precula processed into several peptide hormones MSH, β-and y-LPH, CLIP, β-endorphin, all regulating feeding behavior and POMC-p the target of leptin and insulin for the prorection. 1. Chretien, M. and Mbikay, M. (2016) J M. 2. Yang, Y. et al. (2011) Cell 146, 992-100.	Application Western Blotting Immunoprecipitation Immunofluorescence (Frozen) Immunofluorescence (Immunocytochemistry) Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody POMC (D3R1U) Rabbit mAb recognizes endogenous levels of to Human, Rat, Bovine Monoclonal antibody is produced by immunizing animals with a sresidues surrounding Pro69 of human POMC protein. Pro-opiomelanocortin (POMC) is a precursor protein expressed processed into several peptide hormones and neuropeptides. Ar MSH, β-and γ-LPH, CLIP, β-endorphin, and N-POMC (1). POMC regulating feeding behavior and POMC-producing neurons prom the target of leptin and insulin for the promotion of the browning sinces 1. Chretien, M. and Mbikay, M. (2016) J Mol Endocrinol, T49-62. 2. Yang, Y. et al. (2011) Cell 146, 992-1003.	Application Western Blotting Immunoprecipitation Immunofluorescence (Frozen) Immunofluorescence (Immunocytochemistry) Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glyr 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody. POMC (D3R1U) Rabbit mAb recognizes endogenous levels of total POMC protein. Human, Rat, Bovine Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresidues surrounding Pro69 of human POMC protein. Pro-opiomelanocortin (POMC) is a precursor protein expressed in the pituitary and the processed into several peptide hormones and neuropeptides. Among these peptides MSH, β-and y-LPH, CLIP, β-endorphin, and N-POMC (1). POMC is involved in hypott regulating feeding behavior and POMC-producing neurons promote satiety (2). POMC the target of leptin and insulin for the promotion of the browning of white fat (3). Inces 1. Chretien, M. and Mbikay, M. (2016) J Mol Endocrinol, T49-62. doi: 10.1530/JME-152. Yang, Y. et al. (2011) Cell 146, 992-1003.

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS,

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

WB: Western Blotting IP: Immunoprecipitation IF-F: Immunofluorescence (Frozen) **Applications Key**

IF-IC: Immunofluorescence (Immunocytochemistry)

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster **Cross-Reactivity Key**

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and

Western Blot Buffer

Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the

following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

POMC (D3R1U) Rabbit mAb (#23499) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.