

#66350 Store at -20C

PHGDH (D8F3O) Rabbit mAb**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H	Endogenous	57	Rabbit IgG	#O43175	26227

Product Usage Information**Application**Western Blotting
Immunoprecipitation**Dilution**1:1000
1:50**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

PHGDH (D8F3O) Rabbit mAb recognizes endogenous levels of total PHGDH protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Val304 of human PHGDH protein.

Background

Mammalian cells synthesize serine *de novo* by diverting a portion of the glycolytic intermediate 3-phosphoglycerate into the phosphorylated pathway of serine synthesis. This shift supports anabolism by providing precursors for the biosynthesis of proteins, nucleotides, creatine, porphyrins, phospholipids, and glutathione. Phosphoglycerate dehydrogenase (PHGDH) catalyzes the first step in the serine biosynthesis pathway by converting 3-phosphoglycerate into phosphohydroxy pyruvate (1).

Research studies demonstrate that an increase in serine biosynthesis supports growth and proliferation of cancer cells (2-4), which is supported by amplification and overexpression of PHGDH in a subset of melanoma and breast cancers (5,6). Suppression of PHGDH expression in cell lines with elevated PHGDH levels causes a strong decrease in cell proliferation and inhibits tumor growth *in vivo* (5). Additional evidence suggests that PHGDH interacts with and stabilizes FoxM1, which promotes the proliferation, invasion, and tumorigenicity of glioma cells (7).

Background References

1. Locasale, J.W. (2013) *Nat Rev Cancer* 13, 572-83.
2. Amelio, I. et al. (2013) *Oncogene*, [Epub ahead of print].
3. Ma, L. et al. (2013) *Cell* 152, 599-611.
4. Maddocks, O.D. et al. (2013) *Nature* 493, 542-6.
5. Possemato, R. et al. (2011) *Nature* 476, 346-50.
6. Locasale, J.W. et al. (2011) *Nat Genet* 43, 869-74.
7. Liu, J. et al. (2013) *J Neurooncol* 111, 245-55.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key**WB:** Western Blotting **IP:** Immunoprecipitation**Cross-Reactivity Key**

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
 All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.