e at -20C	ICOS (D1K2T [™]) Rabbit mAb		Cell Signaling	
Store		Orders:	877-616-CELL (2355) orders@cellsignal.com	
01		Support:	877-678-TECH (8324)	
#89601		Web:	info@cellsignal.com cellsignal.com	
Ħ		3 Trask Lane Danvers Ma	ssachusetts 01923 USA	

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: IHC-Bond, IHC-P	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 25-40	Source/Isotype: Rabbit IgG	UniProt ID: #Q9Y6W8	Entrez-Gene Id: 29851		
Product Usage Information		Application		Dilution				
lineination		IC Leica Bond			1:100 - 1:4			
	In	nmunohistochemistry	(Paraffin)		1:100 - 1:4	400		
Storage				7.5), 150 mM NaCl, 100 $\mu g/ml$ BSA, 50% glycerol and less than o not aliquot the antibody.				
		For a carrier free (BSA and azide free) version of this product see product #39740.						
Specificity / Sensitivity		ICOS (D1K2T [™]) Rabbit mAb recognizes endogenous levels of total ICOS protein.						
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human ICOS protein.						
Background		ICOS (Inducible Co-Stimulator, CD278) is a member of the CD28 family that regulates T cell activity and immune responses (1). The ICOS protein contains an extracellular IgV-like domain, a transmembrane domain, and an intracellular domain with a YMFM motif (1-2). ICOS is primarily expressed on activated CD4+ and CD8+ T cells (1). Upon binding to its ligand, ICOS potentiates the T cell response to antigen through activation of the PI3K signaling pathway (3). In addition to enhancing T cell activation and proliferation, ICOS plays an important role in the regulation of T follicular helper cells (4). Research studies suggest that ICOS is a potential therapeutic target, and could serve as a prognostic biomarker for neoplastic therapy involving CTLA-4 blockade (5-7).						
Background References		 Hutloff, A. et al. (1999) Nature 397, 263-6. Coyle, A.J. et al. (2000) Immunity 13, 95-105. Fos, C. et al. (2008) J Immunol 181, 1969-77. Hutloff, A. (2015) Oncotarget 6, 21785-6. Kyi, C. and Postow, M.A. (2014) FEBS Lett 588, 368-76. Fan, X. et al. (2014) J Exp Med 211, 715-25. Ng Tang, D. et al. (2013) Cancer Immunol Res 1, 229-34. 						
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).						
Applications Key		IHC-Bond: IHC Leica Bond IHC-P: Immunohistochemistry (Paraffin)						
Cross-Reactivity Key		 H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected 						
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. SignalStain is a registered trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.				demarks for more		
Limited Uses	follo con	owing terms apply to ditions that are in ad	nerwise expressly agreed in a writing signed by a legally authorized representative of CST, the ns apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and at are in addition to, or different from, those contained herein, unless separately accepted in egally authorized representative of CST, are rejected and are of no force or effect.					

ICOS (D1K2T[™]) Rabbit mAb (#89601) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.