

#97249 Store at -20°C

## ERG (A7L1G) Rabbit mAb



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP, IHC-P, IF-IC, FC-FP	H M	Endogenous	54	Rabbit IgG	#P11308	2078

Product Usage Information	Application	Dilution
	Western Blotting	1:1000
	Immunoprecipitation	1:100
	Immunohistochemistry (Paraffin)	1:200
	Immunofluorescence (Immunocytochemistry)	1:500
	Flow Cytometry (Fixed/Permeabilized)	1:50
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
	For a carrier free (BSA and azide free) version of this product see product #45381.	
Specificity / Sensitivity	ERG (A7L1G) Rabbit mAb recognizes endogenous levels of total ERG protein. Based on sequence identity, this antibody should detect isoforms ERG1, ERG2 and ERG3. This antibody does not cross-react with FLI1.	
Species predicted to react based on 100% sequence homology	Rat, Hamster, Pig, Horse, Guinea Pig	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human ERG protein.	
Background	ETS-related gene (ERG) is a member of the E-26 transformation-specific (ETS) family of sequence-specific DNA-binding transcription factors (1). ERG plays important and highly conserved roles in vertebrate development. Early in embryonic development, ERG is highly expressed in the embryonic mesoderm and endothelium, where it plays a critical role in the formation of the vascular system, urogenital tract and bone development (2,3). Later in embryonic development, ERG functions to regulate the pluripotency of hematopoietic stem cells, endothelial cell homeostasis and angiogenesis (2,4-7). ERG expression is not restricted to development. In adult mouse, ERG is normally expressed in endothelial tissues, including adrenal, cartilage, heart, spleen, lymphatic endothelial and eosinophil cells (8). However, deregulation of ERG activity, often resulting from chromosomal rearrangements, has been implicated and linked to poor prognosis in a number of different cancers. Chromosomal translocations generating EWS/ERG chimeric proteins comprised of the amino-terminal transactivation domain of Ewing's sarcoma breakpoint region 1 (EWS) and the carboxy-terminal ETS domain of ERG have been identified in 5-10% of Ewing's sarcoma, an aggressive bone and soft tissue tumor (9). Chromosomal translocations between ERG and TLS/FUS or ERG and ELF4 have been implicated in acute myeloid leukemia (10, 11). Over-expression of ERG, resulting from gene fusion with the androgen-driven promoter of the TMPRSS2 gene, has been identified as a key driver of metastasis and marker for poor prognosis in prostate cancer (12).	

### Background References

1. Adamo, P. and Ladomery, M.R. (2016) *Oncogene* 35, 403-14.
2. Birdsey, G.M. et al. (2008) *Blood* 111, 3498-506.
3. Vijayaraj, P. et al. (2012) *Development* 139, 3973-85.
4. Ng, A.P. et al. (2011) *Blood* 118, 2454-61.
5. Birdsey, G.M. et al. (2015) *Dev Cell* 32, 82-96.
6. Lathen, C. et al. (2014) *Circulation* 130, 1179-91.
7. McLaughlin, F. et al. (2001) *Blood* 98, 3332-9.
8. Mohamed, A.A. et al. (2010) *J Cancer* 1, 197-208.
9. Chen, S. et al. (2016) *Genes Chromosomes Cancer* 55, 340-9.

10. Ichikawa, H. et al. (1994) *Cancer Res* 54, 2865-8.11. Moore, S.D. et al. (2006) *Leuk Res* 30, 1037-42.12. Tomlins, S.A. et al. (2005) *Science* 310, 644-8.**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key****WB:** Western Blotting **IP:** Immunoprecipitation **IHC-P:** Immunohistochemistry (Paraffin)  
**IF-IC:** Immunofluorescence (Immunocytochemistry) **FC-FP:** Flow Cytometry (Fixed/Permeabilized)**Cross-Reactivity Key****H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected**Trademarks and Patents**Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
Alexa Fluor is a registered trademark of Life Technologies Corporation.  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](https://www.cellsignal.com/trademarks) for more information.**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.