

#58326 Store at +4°C

## Mcl-1 (D2W9E) Rabbit mAb (Alexa Fluor® 488 Conjugate)



**Cell Signaling**  
TECHNOLOGY®

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> FC-FP	<b>Reactivity:</b> H M R	<b>Sensitivity:</b> Endogenous	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #P97287	<b>Entrez-Gene Id:</b> 17210
-------------------------------	-----------------------------	-----------------------------------	--------------------------------------	-------------------------------	---------------------------------

<b>Product Usage Information</b>	<b>Application</b> Flow Cytometry (Fixed/Permeabilized)	<b>Dilution</b> 1:50
<b>Storage</b>	Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.	
<b>Specificity / Sensitivity</b>	Mcl-1 (D2W9E) Rabbit mAb (Alexa Fluor® 488 Conjugate) recognizes endogenous levels of total Mcl-1 protein.	
<b>Source / Purification</b>	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro60 of mouse Mcl-1 protein.	
<b>Product Description</b>	This Cell Signaling Technology antibody is conjugated to Alexa Fluor® 488 fluorescent dye and tested in-house for direct flow cytometric analysis in human cells. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated Mcl-1 (D2W9E) Rabbit mAb #94296.	
<b>Background</b>	Mcl-1 is an anti-apoptotic member of the Bcl-2 family originally isolated from the ML-1 human myeloid leukemia cell line during phorbol ester-induced differentiation along the monocyte/macrophage pathway (1). Similar to other Bcl-2 family members, Mcl-1 localizes to the mitochondria (2), interacts with and antagonizes pro-apoptotic Bcl-2 family members (3), and inhibits apoptosis induced by a number of cytotoxic stimuli (4). Mcl-1 differs from its other family members in its regulation at both the transcriptional and posttranslational level. First, Mcl-1 has an extended amino-terminal PEST region, which is responsible for its relatively short half-life (1,2). Second, unlike other family members, Mcl-1 is rapidly transcribed via a PI3K/Akt dependent pathway, resulting in its increased expression during myeloid differentiation and cytokine stimulation (1,5-7). Mcl-1 is phosphorylated in response to treatment with phorbol ester, microtubule-damaging agents, oxidative stress, and cytokine withdrawal (8-11). Phosphorylation at Thr163, the conserved MAP kinase/ERK site located within the PEST region, slows Mcl-1 protein turnover (10) but may prime the GSK-3 mediated phosphorylation at Ser159 that leads to Mcl-1 destabilization (11). Mcl-1 deficiency in mice results in peri-implantation lethality (12). In addition, conditional disruption of the corresponding <i>mcl-1</i> gene shows that Mcl-1 plays an important role in early lymphoid development and in the maintenance of mature lymphocytes (13).	
<b>Background References</b>	<ol style="list-style-type: none"> <li>1. Kozopas, K.M. et al. (1993) <i>Proc Natl Acad Sci USA</i> 90, 3516-20.</li> <li>2. Yang, T. et al. (1995) <i>J Cell Biol</i> 128, 1173-84.</li> <li>3. Sato, T. et al. (1994) <i>Proc Natl Acad Sci USA</i> 91, 9238-42.</li> <li>4. Zhou, P. et al. (1997) <i>Blood</i> 89, 630-43.</li> <li>5. Wang, J.M. et al. (1999) <i>Mol Cell Biol</i> 19, 6195-206.</li> <li>6. Jourdan, M. et al. (2003) <i>Oncogene</i> 22, 2950-9.</li> <li>7. Chao, J.R. et al. (1998) <i>Mol Cell Biol</i> 18, 4883-98.</li> <li>8. Domina, A.M. et al. (2000) <i>J Biol Chem</i> 275, 21688-94.</li> <li>9. Inoshita, S. et al. (2002) <i>J Biol Chem</i> 277, 43730-4.</li> <li>10. Domina, A.M. et al. (2004) <i>Oncogene</i> 23, 5301-15.</li> <li>11. Maurer, U. et al. (2006) <i>Mol Cell</i> 21, 749-60.</li> <li>12. Rinkenberger, J.L. et al. (2000) <i>Genes Dev</i> 14, 23-7.</li> <li>13. Opferman, J.T. et al. (2003) <i>Nature</i> 426, 671-6.</li> </ol>	

<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
<b>Applications Key</b>	<b>FC-FP:</b> Flow Cytometry (Fixed/Permeabilized)

**Cross-Reactivity Key**

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
Alexa Fluor is a registered trademark of Life Technologies Corporation.  
This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is conditioned on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not (1) use this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; or (c) manufacturing or quality assurance or quality control, and/or (2) sell or transfer this product or its components for resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or [outlicensing@lifetech.com](mailto:outlicensing@lifetech.com).  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.