#88583 Store at -20C

Phospho-HER2/ErbB2 (Thr686) Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:	
WB	Н	Endogenous	185	Rabbit	#P04626	2064	
Product Usage Information	Application			Dilution			
	Western Blotting			1:1000			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity / Sensi		Phospho-HER2/ERBB2 (Thr686) Antibody recognizes endogenous levels of HER2/ERBB2 protein only when phosphorylated at Thr686.					

Species predicted to react based on 100% sequence homology:

Mouse, Rat

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phospho-peptide corresponding to residues surrounding Thr686 of human HER2/ERBB2 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

The ErbB2 (HER2) proto-oncogene encodes a 185 kDa transmembrane, receptor-like glycoprotein with intrinsic tyrosine kinase activity (1). While ErbB2 lacks an identified ligand, ErbB2 kinase activity can be activated in the absence of a ligand when overexpressed and through heteromeric associations with other ErbB family members (2). Amplification of the *ErbB2* gene and overexpression of its product are detected in almost 40% of human breast cancers (3). Binding of the c-Cbl ubiquitin ligase to ErbB2 at Tyr1112 leads to ErbB2 poly-ubiquitination and enhances degradation of this kinase (4). ErbB2 is a key therapeutic target in the treatment of breast cancer and other carcinomas and targeting the regulation of ErbB2 degradation by the c-Cbl-regulated proteolytic pathway is one potential therapeutic strategy. Phosphorylation of the kinase domain residue Tyr877 of ErbB2 (homologous to Tyr416 of pp60c-Src) may be involved in regulating ErbB2 biological activity. The major autophosphorylation sites in ErbB2 are Tyr1248 and Tyr1221/1222; phosphorylation of these sites couples ErbB2 to the Ras-Raf-MAP kinase signal transduction pathway (1,5).

Thr686 is located in the cytoplasmic juxtamembrane region of ErbB2. PKC, PKA, and PKG II kinases have been reported to phosphorylate ErbB2 at this site, but the functional consequences of phosphorylation may be context dependent. For example, one research study reported that phosphorylation of ErbB2 at Thr686 enhanced ligand-induced receptor activation (6), while a separate research study reported inhibitory effects of phosphorylation on ErbB2 activation (8).

Background References

- 1. Muthuswamy, S.K. et al. (1999) Mol Cell Biol 19, 6845-57.
- 2. Qian, X. et al. (1994) Proc Natl Acad Sci USA 91, 1500-4.
- 3. Dittadi, R. and Gion, M. (2000) J Natl Cancer Inst 92, 1443-4.
- 4. Klapper, L.N. et al. (2000) Cancer Res 60, 3384-8.
- 5. Kwon, Y.K. et al. (1997) J Neurosci 17, 8293-9.
- 6. Monje, P.V. et al. (2008) J Biol Chem 283, 34087-100.
- 7. Ouyang, X. et al. (1998) Carcinogenesis 19, 2013-9.
- 8. Zhu, M. et al. (2016) Mol Med Rep 13, 1909-13.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

Cross-Reactivity Key

WB: Western Blotting

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dq: doq Pq: piq Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Limited Uses

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.