

Store at -20C
#2033

DHCR24/Seladin-1 (C59D8) Rabbit mAb



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP, IHC-P	H M	Endogenous	54	Rabbit IgG	#Q15392	1718

Product Usage Information

Application

Western Blotting
Immunoprecipitation
Immunohistochemistry (Paraffin)

Dilution

1:1000
1:50
1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

DHCR24/Seladin-1 (C59D8) Rabbit mAb detects endogenous levels of total DHCR24/Seladin-1 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to the sequence of human DHCR24/Seladin-1.

Background

DHCR24/Seladin-1 was identified as a molecular basis for desmosterolosis (1). It encodes for 24-dehydrocholesterol reductase (3 β -hydroxysterol Δ -24-reductase). This enzyme reduces desmosterol in cholesterol biosynthesis (1). Recessive mutations in this gene in desmosterolosis patients lead to a defective enzyme resulting in increased levels of desmosterol (1). DHCR24/Seladin-1 is induced upon oxidative stress and was found to mediate Ras-induced senescence resulting from increased reactive oxygen species (2). Studies further indicate that the level of DHCR24/Seladin-1 is induced in the acute response and reduced in the chronic response to oxidative stress in a cholesterol dependent manner (3). Moreover, overexpression of DHCR24/Seladin-1 bearing two mutations that abolish its reductase activity causes the cells to lose protection from oxidative stress (3). These findings thus link the reductase activity of DHCR24/Seladin-1 to its protective role in oxidative stress. This enzyme has also been demonstrated to be a hydrogen peroxide scavenger (4).

Background References

1. Waterham, H.R. et al. (2001) *Am J Hum Genet* 69, 685-94.
2. Wu, C. et al. (2004) *Nature* 432, 640-5.
3. Kuehnle, K. et al. (2008) *Mol Cell Biol* 28, 539-50.
4. Lu, X. et al. (2008) *Endocrinology* 149, 3267-73.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation **IHC-P:** Immunohistochemistry (Paraffin)

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.