e at -20C	Axin1 (C95H11) Rabbit mAb		Cell Signaling		
Store at		Orders:	877-616-CELL (2355) orders@cellsignal.com		
74		Support:	877-678-TECH (8324)		
207		Web:	info@cellsignal.com cellsignal.com		
#	3	Trask Lane Danvers	Massachusetts   01923   USA		

Applications	Depativity	Consitivity	B #11A / /
For Research Use On	ly. Not for Use in	Diagnostic Proc	edures.

Applications: WB, IP, IHC-P	Reactivity: H Mk	Sensitivity: Endogenous	<b>MW (kDa):</b> 110	Source/Isotype: Rabbit	<b>UniProt ID:</b> #O15169	Entrez-Gene Id: 8312	
Product Usage	٩	pplication			C	Dilution	
Information	V	Vestern Blotting			1	.:1000	
		mmunoprecipitation				.:50	
	Ir	nmunohistochemistry	(Paraffin)		1	.:50	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.					
Specificity / Sensitivity		Axin1 (C95H11) Rabbit mAb detects the endogenous levels of total Axin1 protein.					
Source / Purification		Monoclonal antibody is produced by immunizing animals with a recombinant protein of human Axin1.					
Background	siq ph is ac wi or pa	gnaling. Axin1 interact nosphorylation and su recruited to the memb stivation of Wnt signali th MEKK1 and activat Axil) can functionally	s with APC, GSk bsequent degrad orane by phosph ing (3,4). In addit tes c-Jun amino- substitute for Axi	2 are multidomain scaff $\zeta$ -3 $\beta$ , Dvl, and $\beta$ -catenin lation of $\beta$ -catenin (1,2). orylated LRP5/6, a proce- ion to its role in the Wnt terminal kinase (JNK/SA in1 in mice (6). Axin2 its the duration and/or inter-	and promotes the GS Upon stimulation of c ess that is believed to signaling pathway, Ax APK) (5). Axin2 (also k elf is a direct target of	K-3β-mediated ells with Wnt, Axin1 be crucial for kin1 forms a complex mown as Conductin the Wnt signaling	
Background Refer	2. 3. 4. 5. 6. 7. 8.	Zeng, X. et al. (2005) Davidson, G. et al. (2 Zhang, Y. et al. (1999	oodgett, J.R. (200 Nature 438, 873 005) Nature 438 ) J. Biol. Chem. 1 tini, F. (2005) Mc ) Mol. Cell Biol. 2 ) Mol. Cell Biol. 2	95) <i>J. Clin. Pathol.</i> 58, 2 8-877. , 867-872. 274, 35247-35254. <i>J. Cell Biol.</i> 25, 4371-43 22, 1172-1183. 22, 1184-1193.			
Species Reactivity		ecies reactivity is dete	ermined by testing	g in at least one approve	ed application (e.g., w	estern blot).	
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				% w/v BSA, 1X TBS,	
Applications Key	w	WB: Western Blotting IP: Immunoprecipitation IHC-P: Immunohistochemistry (Paraffin)					
Cross-Reactivity K	X:	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected					
Trademarks and Patents	U.S All	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.				demarks for more	
Limited Uses							

## Axin1 (C95H11) Rabbit mAb (#2074) Datasheet Without Images Cell Signaling Technology

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.