

**#2141** Store at -20°C

# TANK Antibody


**Cell Signaling**  
TECHNOLOGY®

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

**3 Trask Lane | Danvers | Massachusetts | 01923 | USA**
**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> WB, IP	<b>Reactivity:</b> H M R	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 50	<b>Source:</b> Rabbit	<b>UniProt ID:</b> #Q92844	<b>Entrez-Gene Id:</b> 10010
--------------------------------	-----------------------------	-----------------------------------	------------------------	--------------------------	-------------------------------	---------------------------------

<b>Product Usage Information</b>	<b>Application</b> Western Blotting Immunoprecipitation	<b>Dilution</b> 1:1000 1:50
<b>Storage</b>	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
<b>Specificity / Sensitivity</b>	TANK Antibody detects endogenous levels of total TANK protein. It does not cross react with other TRAF family members.	
<b>Species predicted to react based on 100% sequence homology:</b>	Monkey, Bovine, Dog	
<b>Source / Purification</b>	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ser357 of human TANK. Antibodies are purified by protein A and peptide affinity chromatography.	
<b>Background</b>	<p>TRAFs (TNF receptor-associated factors) are a family of multifunctional adaptor proteins that bind to surface receptors and recruit additional proteins to form multiprotein signaling complexes capable of promoting cellular responses (1-3). Members of the TRAF family share a common carboxy-terminal "TRAF domain", which mediates interactions with associated proteins; many also contain amino-terminal Zinc/RING finger motifs. The first TRAFs identified, TRAF1 and TRAF2, were found by virtue of their interactions with the cytoplasmic domain of TNF-receptor 2 (TNFR2) (4). The six known TRAFs (TRAF1-6) act as adaptor proteins for a wide range of cell surface receptors and participate in the regulation of cell survival, proliferation, differentiation, and stress responses.</p> <p>TRAF-associated NF-κB activator (TANK), also known as TRAF-interacting protein (I-TRAF), is a TRAF binding protein that demonstrates both stimulatory and inhibitory properties (5,6). TANK binds to the carboxy domain of the TRAF1, -2 and -3. Overexpression of TANK prevents the association of TRAF2 with TNFR2, inhibiting TNFR2 and CD40 induced NF-κB activation (6). TANK is also reported to synergize with low levels of TRAF2 to activate NF-κB (5). TANK assists in the activation of NF-κB via association and activation of TANK-binding kinase 1 (TBK1) or IKKε which promotes activation of the IKK complex (7,8). It has also been shown that TANK may synergize with TRAF2, TRAF5, and TRAF6 but not TRAF3 in SAPK activation (9). TNFα stimulation results in IKKβ-dependent phosphorylation of TANK which may provide negative feedback regulation of TANK mediated NF-κB activation (10).</p>	
<b>Background References</b>	<ol style="list-style-type: none"> <li>1. Arch, R.H. et al. (1998) <i>Genes Dev</i> 12, 2821-30.</li> <li>2. Chung, J.Y. et al. (2002) <i>J Cell Sci</i> 115, 679-88.</li> <li>3. Bradley, J.R. and Pober, J.S. (2001) <i>Oncogene</i> 20, 6482-91.</li> <li>4. Rothe, M. et al. (1994) <i>Cell</i> 78, 681-92.</li> <li>5. Cheng, G. and Baltimore, D. (1996) <i>Genes Dev.</i> 10, 963-973.</li> <li>6. Rothe, M. et al. (1996) <i>Proc. Natl. Acad. Sci. USA</i> 93, 8241-8846.</li> <li>7. Pomerantz, J.L. and Baltimore, D. (1999) <i>EMBO J.</i> 18, 6694-6704.</li> <li>8. Chariot, A. et al. (2002) <i>J. Biol. Chem.</i> 277, 37029-37036.</li> <li>9. Chin, A.I. et al. (1999) <i>Mol. Cell. Biol.</i> 19, 6665-6672.</li> <li>10. Bonif, M. et al. (2006) <i>Biochem. J.</i> 394, 593-603.</li> </ol>	

## Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**WB:** Western Blotting **IP:** Immunoprecipitation

**Cross-Reactivity Key**

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](https://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.