Revision 3							
Phospho-Chk1 (Ser345) (133D3) Rabbit mAb					Cell Signaling TECHNOLOGY* Orders: 877-616-CELL (2355)		
					Support	orders@cellsignal.com 877-678-TECH (8324)	
48					Support:		
#2348					Web:	info@cellsignal.com cellsignal.com	
				3 Trask I	_ane Danvers Ma	ssachusetts 01923 USA	
For Research Use Onl	•						
Applications: WB, IF-IC, FC-FP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 56	Source/Isotype: Rabbit IgG	UniProt ID: #014757	Entrez-Gene Id: 1111	
Product Usage	Δ	pplication				Dilution	
Information		/estern Blotting				1:1000	
		nmunofluorescence (Immunocytochem	nistry)		1:50	
		ow Cytometry (Fixed	. ,	,		1:200	
Storage	Su	pplied in 10 mM sod	ium HEPES (pH 7	7.5), 150 mM NaCl, 100 not aliquot the antibod		ycerol and less than	
	Fo	r a carrier free (BSA	and azide free) ve	ersion of this product se	e product #76784.		
Specificity / Sensitivity		Phospho-Chk1 (Ser345) (133D3) Rabbit mAb detects endogenous levels of Chk1 only when phosphorylated at serine 345.					
Source / Purificat		Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser345 of human Chk1.					
Background Background References		 Chk1 kinase acts downstream of ATM/ATR kinase and plays an important role in DNA damage checkpoint control, embryonic development, and tumor suppression (1). Activation of Chk1 involves phosphorylation at Ser317 and Ser345 by ATM/ATR, followed by autophosphorylation of Ser296. Activation occurs in response to blocked DNA replication and certain forms of genotoxic stress (2). While phosphorylation at Ser345 serves to localize Chk1 to the nucleus following checkpoint activation (3), phosphorylation at Ser317 along with site-specific phosphorylation of PTEN allows for re-entry into the cell cycle following stalled DNA replication (4). Chk1 exerts its checkpoint mechanism on the cell cycle, in part, by regulating the cdc25 family of phosphatases. Chk1 phosphorylation of cdc25A targets if for proteolysis and inhibits its activity through 14-3-3 binding (5). Activated Chk1 can inactivate cdc25C via phosphorylation at Ser216, blocking the activation of cdc2 and transition into mitosis (6). Centrosomal Chk1 has been shown to phosphorylate cdc25B and inhibit its activation of CDK1-cyclin B1, thereby abrogating mitotic spindle formation and chromatin condensation (7). Furthermore, Chk1 plays a role in spindle checkpoint function through regulation of aurora B and BubR1 (8). Research studies have implicated Chk1 as a drug target for cancer therapy as its inhibition leads to cell death in many cancer cell lines (9). 1. Liu, Q. et al. (2000) <i>Genes Dev</i> 14, 1448-59. 2. Zhao, H. and Piwnica-Worms, H. (2001) <i>Mol Cell Biol</i> 21, 4129-39. 3. Jiang, K. et al. (2003) <i>J Biol Chem</i> 278, 25207-17. 4. Martin, S.A. and Ouchi, T. (2008) <i>Mol Cancer Ther</i> 7, 2509-16. 5. Chen, M.S. et al. (2003) <i>Nol Cell Biol</i> 23, 7488-97. 6. Zeng, Y. et al. (1998) <i>Nature</i> 395, 507-10. 7. Löffler, H. et al. (2006) <i>Cell Cycle</i> 5, 2543-7. 					
Western Blot Buffer IMPORTANT: For western blo 0.1% Tween® 20 at 4°C with			Vatl Cancer Inst 9 ermined by testing rn blots, incubate with gentle shaki	Cancer Inst 97, 1026-8. ned by testing in at least one approved application (e.g., western blot). ots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, a gentle shaking, overnight. : Immunofluorescence (Immunocytochemistry)			
Cross-Reactivity	FC	-FP: Flow Cytometry	/ (Fixed/Permeabi	lized)			

1/1/24, 12:01 PM	Phospho-Chk1 (Ser345) (133D3) Rabbit mAb (#2348) Datasheet Without Images Cell Signaling Technology H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.
	Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.