e at -20C	Rab11b Antibody		Cell Signaling		
Store at		Orders:	877-616-CELL (2355) orders@cellsignal.com		
[4		Support:	877-678-TECH (8324)		
#2414		Web:	info@cellsignal.com cellsignal.com		
#		3 Trask Lane Danvers	Massachusetts 01923 USA		

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H M R B	Sensitivity: Endogenous	MW (kDa): 25	Source: Rabbit	UniProt ID: #Q15907	Entrez-Gene Id: 9230		
Product Usage Information		oplication estern Blotting			Dilution 1:1000			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.						
Specificity / Sensitivity		Rab11b Antibody detects endogenous levels of total Rab11b protein.						
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding residue surrounding Arg184 of human Rab11b protein. Antibodies are purified by protein A and peptide affinity chromatography.						
Background		Rab11a, Rab11b and Rab25 are members of the Rab11 family of small Ras-like GTPases. Rab11 (isoforms Rab11a and Rab11b) functions as a key regulator in the recycling of perinuclear, plasma membrane and Golgi compartment endosomes (1,2). Despite some overlap, distinct differences exist between Rab11a and Rab11b in both their cellular distribution and functional roles. Rab11a is ubiquitously expressed while Rab11b is found mainly in the heart and brain (3,4). Like other Rab proteins, Rab11 exerts its function via interactions with Rab11 family interacting proteins (FIPs). While there are three distinct classes of FIPs, all appear to share a conserved carboxy-terminal Rab-binding domain that allows Rab-FIP protein interaction. When bound together, these proteins are thought to regulate membrane-associated protein sorting (5,6).						
Background Refe	2. C 3. L 4. k 5. J	Jllrich, O. et al. (1996 Chen, W. et al. (1998) apierre, L.A. et al. (2 Chvotchev, M.V. et al. unutula, J.R. et al. (20 Iales, C.M. et al. (200	Mol. Biol. Cell 9, 3 003) Exp. Cell Res (2003) J. Neurosc 2004) J. Biol. Chem	3241-3257. 5. 290, 322-331. 6. 23, 10531-10539. 5. 279, 33430-33437				
Species Reactivit	y Spe	Species reactivity is determined by testing in at least one approved application (e.g., western blot).						
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications Key	WB	: Western Blotting						
Cross-Reactivity Key		H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected						
Trademarks and Patents	All o	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.						
Limited Uses	follo conc	wing terms apply to F ditions that are in add	otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the erms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and that are in addition to, or different from, those contained herein, unless separately accepted in a legally authorized representative of CST, are rejected and are of no force or effect.					

Rab11b Antibody (#2414) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.