

#2817 Store at -20C

MLK3 Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M R Hm Mk	Endogenous	92	Rabbit	#Q16584	4296

Product Usage Information	Application Western Blotting	Dilution 1:1000
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	MLK3 Antibody detects endogenous levels of total MLK3 protein. This antibody does not cross-react with MLK1, MLK2 and other mixed lineage kinases.	
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human MLK3. Antibodies are purified by protein A and peptide affinity chromatography.	
Background	Mixed lineage kinase 3 (MLK3) is a serine/threonine kinase that has an amino-terminal SH3 domain followed by the kinase domain and two leucine zippers, a cdc42/Rac1 binding (CRIB) domain and several other domains/motifs at the carboxy-terminal region. CRIB triggers the dimerization of MLK3 via its tandem leucine zippers, followed by the intramolecular phosphorylation and subsequent activation of MLK3 (1,2). Autophosphorylation of Thr277 and Ser281 is essential for MLK3 kinase activity (3). Ser281 is also phosphorylated by HPK in an <i>in vitro</i> kinase assay (3). MLK3 functions as a MAPKKK of the SAPK/JNK stress pathway by directly phosphorylating SEK1/MKK4 and MKK7, although it is controversial whether MLK3 is involved in p38 stress pathway activation (1,4). MLK3 also functions as an IκB kinase and mediates the activation of the transcriptional factor NF-κB stimulated by CD3/CD28, suggesting a role for MLK3 in immune and inflammatory responses (5).	
Background References	<ol style="list-style-type: none"> 1. Teramoto, H. et al. (1996) <i>J. Biol. Chem.</i> 271, 27225-27228. 2. Leung, I. W. et al. (1998) <i>J. Biol. Chem.</i> 273, 32408-32415. 3. Leung, I.W. and Lassam, N. (2001) <i>J. Biol. Chem.</i> 276, 1961-1967. 4. Tibbles, L. A. et al. (1996) <i>EMBO J.</i> 15, 7026-7035. 5. Hehner, S. P. et al. (2000) <i>Mol. Cell. Biol.</i> 20, 2556-2568. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.