CHOP (L63F7) Mouse mAb				Cell Signaling	
Stor				Orders:	877-616-CELL (2355) orders@cellsignal.com
				Support:	877-678-TECH (8324)
#2895				Web:	info@cellsignal.com cellsignal.com
3 Trask Lane   Danvers   Massachusetts   01923   USA   For Research Use Only. Not for Use in Diagnostic Procedures.					
			0		
Applications: Reactiv WB, IP, IF-IC, FC-FP, H M F ChIP		<b>MW (kDa):</b> 27	Source/Isotype: Mouse IgG2a	UniProt ID: #P35638	Entrez-Gene Id: 1649
Product Usage Information	For optimal ChIP results, use 2.5 $\mu$ l of antibody and 10 $\mu$ g of chromatin (approximately 4 x 10 <sup>6</sup> cells) per IP. This antibody has been validated using SimpleChIP <sup>®</sup> Enzymatic Chromatin IP Kits.				
Application			Dilution		
	Western Blotting			1:1000	
	Immunoprecipitation			1:50	
	Immunofluorescence (Immunocytochemistry)			1:400 - 1:1600	
	Flow Cytometry (Fixed/Permeabilized)			1:250 - 1:1000	
	Chromatin IP			1:	200
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
For a carrier free (BSA and azide free) version of this product see product #78063.					
Specificity / Sensitivity	CHOP (L63F7) Mouse mAb detects endogenous levels of total CHOP protein.				
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to the sequence of human CHOP.				
Background	CHOP was identified as a C/EBP-homologous protein that inhibits C/EBP and LAP in a dominant-negative manner (1). CHOP expression is induced by certain cellular stresses including starvation and the induced CHOP suppresses cell cycle progression from G1 to S phase (2). Later it was shown that, during ER stress, the level of CHOP expression is elevated and CHOP functions to mediate programmed cell death (3). Studies also found that CHOP mediates the activation of GADD34 and Ero1-L $\alpha$ expression during ER stress. GADD34 in turn dephosphorylates phospho-Ser51 of elF2 $\alpha$ thereby stimulating protein synthesis. Ero1-L $\alpha$ promotes oxidative stress inside the endoplasmic reticulum (ER) (4). The role of CHOP in the programmed cell death of ER-stressed cells is correlated with its role promoting protein synthesis and oxidative stress inside the ER (4).				
Background References	1. Ron, D. and Habener 2. Barone, M.V. et al. (1 3. Zinszner, H. et al. (19 4. Marciniak, S.J. et al.	994) Genes Dev 198) Genes Dev 1	8, 453-64. L2, 982-95.		
Species Reactivity	Species reactivity is dete	ermined by testing	g in at least one approve	ed application (e.g.	, western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key	WB: Western Blotting IP: Immunoprecipitation IF-IC: Immunofluorescence (Immunocytochemistry) FC-FP: Flow Cytometry (Fixed/Permeabilized) ChIP: Chromatin IP				
Cross-Reactivity Key	H: human M: mouse R: X: Xenopus Z: zebrafish GP: Guinea Pig Rab: ral	B: bovine Dg: de	og <b>Pg:</b> pig <b>Sc:</b> S. cerevi		

1/1/24, 11:22 AM

Trademarks and Patents

Limited Uses

CHOP (L63F7) Mouse mAb (#2895) Datasheet Without Images Cell Signaling Technology

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.