2909 Store at -20C

## Phospho-(Ser/Thr) ATM/ATR Substrate (4F7) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Dilution

1:1000

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:

Reactivity:

Sensitivity: Endogenous Source/Isotype: Rabbit IgG

Product Usage Information

**Application**Western Blotting

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100  $\mu$ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at  $-20^{\circ}$ C. Do not aliquot the antibody.

Specificity / Sensitivity

Phospho-(Ser/Thr) ATM/ATR Substrate (4F7) Rabbit mAb detects endogenous levels of proteins containing the ATM/ATR substrate motif. This antibody preferentially binds peptides and proteins that contain phospho-Ser/Thr preceded by Leu or similar hydrophobic amino acids at the -1 position and followed by Gln at the +1 position. The antibody does not cross-react with corresponding nonphosphorylated sequences or with other phospho-Ser/Thr-containing motifs. (U.S. Patent No's.:

nonphosphorylated sequences or with other phospho-Ser/Thr-containing motifs. (U.S. Patent No's 6,441,140; 6,982,318; 7,259,022; 7,344,714; U.S.S.N. 11,484,485; and all foreign equivalents.)

Source / Purification

Monoclonal antibody is produced by immunizing animals with synthetic phospho-ATM/ATR substrate peptides.

**Background** 

Ataxia telangiectasia mutated kinase (ATM) and ataxia telangiectasia and Rad3-related kinase (ATR) are related kinases that regulate cell cycle checkpoints and DNA repair (1). The identified substrates for ATM are p53, p95/NBS1, MDM2, Chk2, BRCA1, CtIP, 4E-BP1, and Chk1 (1,2) The essential requirement for the substrates of ATM/ATR is S\*/T\*Q. Hydrophobic amino acids at positions -3 and -1, and negatively charged amino acids at position +1 are positive determinants for substrate recognition by these kinases. Positively charged residues surrounding the S\*/T\*Q are negative determinants for substrate phosphorylation (3). The complex phenotype of AT cells suggests that it likely has additional substrates (3). To better understand the kinase and identify substrates for ATM and the related kinase ATR, CST has developed antibodies that recognize phosphorylated serine or threonine in the S\*/T\*O motif.

## **Background References**

- 1. Kastan, M.B. and Lim, D.S. (2000) Nature Rev. Mol. Cell Biol. 1, 179-186.
- 2. Zhao, H. and Piwnica-Worms, H. (2001) Mol. Cell. Biol. 21, 4129-4139.
- 3. Kim, S. T. et al. (1999) J. Biol. Chem. 274, 37538-37543.

**Species Reactivity** 

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer** 

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key** 

WB: Western Blotting

**Cross-Reactivity Key** 

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

 $\hbox{Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. } \\$ 

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

nformation.

**Limited Uses** 

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

1/1/24. 9:35 AM

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.