

#2996 Store at -20°C

SET8 (C18B7) Rabbit mAb


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IF-IC	H M R Mk	Endogenous	43	Rabbit IgG	#Q9NQR1	387893

Product Usage Information	Application Western Blotting Immunofluorescence (Immunocytochemistry)	Dilution 1:1000 1:100
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	SET8 (C18B7) Rabbit mAb detects endogenous levels of total SET8 protein (both isoforms). The antibody also crossreacts with an unidentified protein at 50kDa.	
Species predicted to react based on 100% sequence homology:	Bovine, Pig, Horse	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to human SET8 protein. Antibodies are purified by protein A and peptide affinity chromatography.	
Background	SET domain-containing lysine methyltransferase 8 (SET8), also known as PR/SET domain-containing protein 7 (PR/SET7), is a member of a family of histone lysine methyltransferases, each of which contains a conserved catalytic SET domain originally identified in <i>Drosophila</i> Su[<i>var</i>]3-9, Enhancer of zeste, and Trithorax proteins (1-3). SET8 is a single-subunit enzyme that mono-methylates histone H4 on Lys20, preferably on nucleosomal substrates (1-3). SET8 protein levels and Histone H4 Lys20 methylation are cell cycle regulated, both increasing in S phase and peaking at G2/M phase (4,5). SET8 interacts with the PCNA protein, associates with sites of active DNA synthesis, and is required for DNA replication and genome stability during S phase (5-7). Inhibition of SET8 using shRNA or siRNA results in arrest of replication forks, induction of double-stranded DNA breaks, and a Chk1-mediated cell-cycle arrest in S and G2/M phases of the cell cycle (6,7). Furthermore, SET8 methylates p53 on Lys382, down regulating the pro-apoptotic and checkpoint activation functions of p53 (8). In response to DNA damage, SET8 expression levels decrease, allowing p53 to activate checkpoints and/or apoptosis (8). Both the methylation of histone H4 Lys20 and p53 appear to be important for the functions of SET8 in S phase.	
Background References	1. Fang, J. et al. (2002) <i>Curr Biol</i> 12, 1086-99. 2. Xiao, B. et al. (2005) <i>Genes Dev</i> 19, 1444-54. 3. Couture, J.F. et al. (2005) <i>Genes Dev</i> 19, 1455-65. 4. Rice, J.C. et al. (2002) <i>Genes Dev</i> 16, 2225-30. 5. Huen, M.S. et al. (2008) <i>J Biol Chem</i> 283, 11073-7. 6. Tardat, M. et al. (2007) <i>J Cell Biol</i> 179, 1413-26. 7. Jørgensen, S. et al. (2007) <i>J Cell Biol</i> 179, 1337-45. 8. Shi, X. et al. (2007) <i>Mol Cell</i> 27, 636-46.	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting IF-IC: Immunofluorescence (Immunocytochemistry)
Cross-Reactivity Key	

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.