36 Store at -200

S5a/PSMD4 (D17E4) Rabbit mAb



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support:

877-678-TECH (8324)

Web:

info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 50	Source/Isotype: Rabbit IgG	UniProt ID: #P55036	Entrez-Gene Id: 5710	
Product Usage Information	•	Application Western Blotting			Dilution 1:1000		
		nunoprecipitation			1:1000		
Storage	•	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.					
Specificity / Sens	itivity S5a	S5a/PSMD4 (D17E4) Rabbit mAb recognizes endogenous levels of total S5a/PSMD4 protein.					
Species predicted react based on 10 sequence homological contractions are sequenced to the contraction of th	00%	nster, Xenopus, Zeb	orafish, Bovine, D	og, Pig, Horse			
Source / Purificat	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to esidues near the amino terminus of human S5a/PSMD4 protein.						
Background	S5a (PSMD4) is a subunit of the 19S regulatory proteasome complex functioning in ubiquitinated-targeting and degradation (1). S5a contains two polyubiquitin binding motifs (UIM) that bind multiu chains by hydrophobic interaction (2,3). In addition to ubiquitin, the UIM of S5a shows high affinity ubiquitin-like domain present in many proteins. S5a binds to these types of proteins directly and me their targeting to the proteasome for degradation (4,5).				t bind multiubiquitin high affinity to a		
Background Refe	2. Yo 3. B 4. F	 Deveraux, Q. et al. (1995) J. Biol. Chem. 270, 23726-23729. Young, P. et al. (1998) J. Biol. Chem. 273, 5461-5467. Beal, R. et al. (1996) Proc. Natl. Acad. Sci. USA 93, 861-866. Fujiwara, K. et al. (2004) J. Biol. Chem. 279, 4760-4767. Sakata, E. et al. (2003) EMBO Rep. 4, 301-306. 					

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dq: doq Pq: piq Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

S5a/PSMD4 (D17E4) Rabbit mAb (#3336) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.