

#3429 Store at -20°C

Phospho-NuMA (Ser395) Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H	Endogenous	238	Rabbit	#Q14980	4926

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Phospho-NuMA (Ser395) Antibody detects endogenous levels of NuMA protein only when phosphorylated at Ser395.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser395 of human NuMA. Antibodies are purified using protein A and peptide affinity chromatography.

Background

The nuclear mitotic apparatus protein (NuMA) is a coiled coil protein involved in the formation and maintenance of the mitotic spindle. NuMA plays a role in chromatin organization during interphase, which influences mammary epithelial differentiation (1,2). During apoptosis, carboxy-terminal cleavage of NuMA may amplify signaling in the cell death pathway (2). NuMA is phosphorylated at numerous sites, with phosphorylation at Ser395 occurring in an ATM/ATR-dependent manner in response to DNA damage (3,4). Phosphorylation at Thr2055 by CDK1 is required for spindle pole association of NuMA at the onset of mitosis. Dephosphorylation by PPP2CA leads to enhancement of NuMA at the cell cortex in anaphase and proper cell-cycle progression (5,6).
Phospho-NuMA (Ser395) Antibody is directed at a site that was identified at Cell Signaling Technology (CST) using PhosphoScan®, CST's LC-MS/MS platform for modification site discovery. Phosphorylation at Ser395 was discovered using an ATM/ATR substrate antibody and was shown to be induced by UV treatment. Please visit PhosphoSitePlus®, CST's modification site knowledgebase, at www.phosphosite.org for more information.

Background References

1. Abad, P.C. et al. (2007) *Mol Biol Cell* 18, 348-61.
2. Lin, H.H. et al. (2007) *J Biomed Sci* 14, 681-94.
3. Stokes, M.P. et al. (2007) *Proc Natl Acad Sci USA* 104, 19855-60.
4. Matsuoka, S. et al. (2007) *Science* 316, 1160-6.
5. Kotak, S. et al. (2013) *EMBO J* 32, 2517-29.
6. Seldin, L. et al. (2013) *Mol Biol Cell* 24, 3651-62.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.