

#3451 Store at -20°C

## Phospho- $\mu$ -Opioid Receptor (Ser375) Antibody



**Cell Signaling**  
TECHNOLOGY®

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	M	Transfected Only	70 to 90	Rabbit	#P35372	4988

### Product Usage Information

#### Application

Western Blotting  
Immunoprecipitation

#### Dilution

1:1000  
1:100

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100  $\mu$ g/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

### Specificity / Sensitivity

Phospho- $\mu$ -Opioid Receptor (Ser375) Antibody detects transfected  $\mu$ -opioid receptor only when phosphorylated at Ser375 of mouse MOR (or Ser377 of human MOR).

### Species predicted to react based on 100% sequence homology:

Human

### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser377 of human (homologous to Ser375 of mouse)  $\mu$ -opioid receptor. Antibodies are purified by protein A and peptide affinity chromatography.

### Background

The  $\mu$ -opioid receptor (MOR) belongs to the superfamily of G-protein-coupled receptors. MOR mediates the analgesic and rewarding effects of morphine and other opiates as well as the actions of several endogenous opioid peptides (1). Upon binding to its ligands, this Gi-coupled receptor inactivates adenylyl cyclase (1) and activates a variety of G-beta-gamma-dependent pathways including the MAPK and the PI3K/Akt cascades (2,3). Trafficking of these receptors to and from the plasma membrane and their desensitization play a significant role in morphine tolerance (4,5). As with other GPCRs, these processes are modulated by phosphorylation at diverse sites within intracellular domains (6). Among other sites, agonist-specific phosphorylation of serine 375 in mouse (serine 377 in human) MOR is essential for its internalization (7).

### Background References

1. Law, P. Y. et al. (2000) *Annu. Rev. Pharmacol. Toxicol.* 40, 389-430.
2. Polakiewicz, R. D. et al. (1998) *J. Biol. Chem.* 273, 12402-12406.
3. Polakiewicz, R. D. et al. (1998) *J. Biol. Chem.* 273, 23534-23541.
4. Finn, A.K. and Whistler, J.L. (2001) *Neuron* 32, 829-839.
5. Kieffer, B.L. and Evans, C.J. (2002) *Cell* 108, 587-590.
6. Yu, Y. et al. (1997) *J. Biol. Chem.* 272, 28869-28874.
7. El Kouhen, R. et al. (2001) *J. Biol. Chem.* 276, 12774-12780.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**WB:** Western Blotting **IP:** Immunoprecipitation

### Cross-Reactivity Key

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](https://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.