#3463 Store at -20C

Phospho-FLT3 (Tyr969) (C24D9) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H M	Sensitivity: Endogenous	MW (kDa): 160	Source/Isotype: Rabbit IgG	UniProt ID: #P36888	Entrez-Gene Id: 2322	
Product Usage Information		Application Western Blotting		Dilution 1:1000			
Storage		•	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	7.5), 150 mM NaCl, 100 o not aliquot the antibody	10 , 0)	cerol and less than	
Specificity / Sensitivity		Phospho-FLT3 (Tyr969) (C24D9) Rabbit mAb detects endogenous levels of FLT3 only when phosphorylated at Tyr969. The antibody may cross-react with other tyrosine-phosphorylated proteins. (US Patent No. 7,183,385 and foreign equivalents assigned to Cell Signaling Technology, Inc.)					
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to the sequence surrounding Tyr969 of human FLT3.					
Background	kina hem (1,2 auto kina Tyr5 FLT	FMS-related tyrosine kinase 3 (FLT3, also called FLK2) is a member of the Type III receptor tyrosine kinase family, which includes c-Kit, PDGFR, and M-CSF receptors. FLT3 is expressed on early hematopoietic progenitor cells and supports growth and differentiation within the hematopoietic system (1,2). FLT3 is activated after binding with its ligand FL, which results in a cascade of tyrosine autophosphorylation and tyrosine phosphorylation of downstream substrates (3). The p85 subunit of PI3 kinase, SHP2, GRB2, and Shc have all been reported to associate with FLT3 after FL stimulation (4-6). Tyr589/591 is located in the juxtamembrane region of FLT3 and may play an important role in regulation of FLT3 tyrosine kinase activity. Somatic mutations of FLT3 consisting of internal tandem duplications (ITDs) occur in 20% of patients with acute myeloid leukemia (7).					
	pho: Pho	Phospho-FLT3 (Tyr969) (C24D9) Rabbit mAb is directed against a previously unpublished FLT3 tyrosine phosphorylation site at Tyr969 that was identified at Cell Signaling Technology (CST) using PhosphoScan [®] , CST's MS/MS platform for phosphorylation site discovery. Phosphorylation of FLT3 at Tyr969 was observed in various leukemia, lymphoma and carcinoma cell lines and in tumors.					
 Shurin, M.R. et al. (1998) Cytokine Growth Factor Rev 9, 37-48. Naoe, T. et al. (2001) Cancer Chemother Pharmacol 48 Suppl 1, S27-30. Namikawa, R. et al. (1996) Stem Cells 14, 388-95. Beslu, N. et al. (1996) J Biol Chem 271, 20075-81. Zhang, S. and Broxmeyer, H.E. (2000) Biochem Biophys Res Commun 277, 195-9. Zhang, S. et al. (1999) J Leukoc Biol 65, 372-80. Mizuki, M. et al. (2000) Blood 96, 3907-14. 							

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

1/1/24. 8:49 AM

Phospho-FLT3 (Tyr969) (C24D9) Rabbit mAb (#3463) Datasheet Without Images Cell Signaling Technology

Trademarks and Patents

Limited Uses

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.