LDHA (C4B5) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Applications: WB, IHC-P, IF-IC, FC- FP	Reactivity: H Mk	Sensitivity: Endogenous	MW (kDa): 37	Source/Isotype: Rabbit IgG	UniProt ID: #P00338	Entrez-Gene Id 3939	
Product Usage Information	Ар	Application			Dilution		
	We	Western Blotting				1:1000	
	Imi	Immunohistochemistry (Paraffin)				1:200 - 1:800	
	Imi	Immunofluorescence (Immunocytochemistry)				1:100 - 1:400	
	Flo	Flow Cytometry (Fixed/Permeabilized)				1:50 - 1:200	
Specificity / Sensi	0.02 For tivity LDH	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody. For a carrier free (BSA and azide free) version of this product see product #43723. LDHA (C4B5) Rabbit mAb detects endogenous levels of total LDHA protein.					
Source / Purificati	•	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to the sequence of human LDHA.					
Background	Who gen the und mus	Lactate dehydrogenase (LDH) catalyzes the interconversion of pyruvate and NADH to lactate and NAD ⁺ . When the oxygen supply is too low for mitochondrial ATP production, this reaction recycles NADH generated in glycolysis to NAD ⁺ , which reenters glycolysis. The major form of LDH found in muscle cells is the A (LDHA) isozyme. The LDHA promoter contains HIF-1α binding sites (1). LDHA expression is induced under hypoxic conditions (2). During intensive and prolonged muscle exercise, lactate accumulates in muscle cells when the supply of oxygen does not meet demand. When oxygen levels return to normal, LDH converts lactate to pyruvate to generate ATP in the mitochondrial electron transport chain.					
Background Refer	rences 1. S	1. Semenza, G.L. et al. (1996) <i>J Biol Chem</i> 271, 32529-37.					

2. Semenza, G.L. (2007) Biochem J 405, 1-9.

Species reactivity is determined by testing in at least one approved application (e.g., western blot). **Species Reactivity**

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS,

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key WB: Western Blotting IHC-P: Immunohistochemistry (Paraffin)

IF-IC: Immunofluorescence (Immunocytochemistry) FC-FP: Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the **Limited Uses**

following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

LDHA (C4B5) Rabbit mAb (#3582) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.