

#3611 Store at -20°C

Phospho-Tuberin/TSC2 (Thr1462) Antibody


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H M	Sensitivity: Endogenous	MW (kDa): 200	Source: Rabbit	UniProt ID: #P49815	Entrez-Gene Id: 7249
----------------------------	---------------------------	-----------------------------------	-------------------------	--------------------------	-------------------------------	--------------------------------

Product Usage Information	Application Western Blotting	Dilution 1:1000
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	Phospho-Tuberin/TSC2 (Thr1462) Antibody detects endogenous levels of tuberin only when phosphorylated at threonine 1462. This antibody does not detect tuberin phosphorylated at other sites.	
Species predicted to react based on 100% sequence homology:	Rat	
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Thr1462 of human tuberin. Antibodies are purified by protein A and peptide affinity chromatography.	
Background	<p>Tuberin is a product of the TSC2 tumor suppressor gene and an important regulator of cell proliferation and tumor development (1). Mutations in either <i>TSC2</i> or the related <i>TSC1</i> (hamartin) gene cause tuberous sclerosis complex (TSC), an autosomal dominant disorder characterized by development of multiple, widespread non-malignant tumors (2). Tuberin is directly phosphorylated at Thr1462 by Akt/PKB (3). Phosphorylation at Thr1462 and Tyr1571 regulates tuberin-hamartin complexes and tuberin activity (3-5). In addition, tuberin inhibits the mammalian target of rapamycin (mTOR), which promotes inhibition of p70 S6 kinase, activation of eukaryotic initiation factor 4E binding protein 1 (4E-BP1, an inhibitor of translation initiation), and eventual inhibition of translation (3,6,7).</p> <p>Tuberin is phosphorylated on Ser939 and Thr1462 in response to PI3K activation, and that the human TSC complex is a direct biochemical target of the PI3K/Akt pathway (3). This data complements Drosophila genetics studies suggesting the possible involvement of the tuberin-hamartin complex in the PI3K/Akt mediated insulin pathway (8-10).</p>	
Background References	<ol style="list-style-type: none"> 1. Soucek, T. et al. (1998) <i>Proc Natl Acad Sci U S A</i> 95, 15653-8. 2. Sparagana, S.P. and Roach, E.S. (2000) <i>Curr Opin Neurol</i> 13, 115-9. 3. Manning, B.D. et al. (2002) <i>Mol Cell</i> 10, 151-62. 4. Aicher, L.D. et al. (2001) <i>J Biol Chem</i> 276, 21017-21. 5. Dan, H.C. et al. (2002) <i>J Biol Chem</i> 277, 35364-70. 6. Goncharova, E.A. et al. (2002) <i>J Biol Chem</i> 277, 30958-67. 7. Inoki, K. et al. (2002) <i>Nat Cell Biol</i> 4, 648-57. 8. Gao, X. and Pan, D. (2001) <i>Genes Dev.</i> 15, 1383-1392. 9. Potter, C. J. et al. (2001) <i>Cell</i> 105, 357-368. 10. Tapon, N. et al. (2001) <i>Cell</i> 105, 345-355. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting
Cross-Reactivity Key	

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.