3672 Store at -200

Myosin Light Chain 2 Antibody



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support:

877-678-TECH (8324)

Web:

info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

	Lanc Danvers Inas	aciiusciis 01323 03A				
For Research Use Only. Not for Use in Diagnostic Procedures.						
Applications: WB	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 18	Source: Rabbit	UniProt ID: #P24844	Entrez-Gene Id: 10398
Product Usage	Ар	plication			Dilution	
Information	We	estern Blotting		1:1000		
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity / Sen	This	Myosin Light Chain 2 Antibody detects endogenous levels of total Myosin Light Chain 2 (smooth muscle). This antibody does not cross-react with the cardiac isoform of Myosin Light Chain 2, or Myosin Essential Light Chain.				
Species predicte react based on 1 sequence homo	.00%	cken, Bovine, Pig				
Source / Purifica	amii	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to amino-terminal residues of human Myosin Light Chain 2 (smooth muscle isoform). Antibodies are purified by protein A and peptide affinity chromatography.				
Background	Myosin is composed of six polypeptide chains: two identical heavy chains and two pairs of light chains. Myosin light chain 2 (MLC2), also known as myosin regulatory light chain (MRLC), RLC, or LC20, has					

many isoforms depending on its distribution. In smooth muscle, MLC2 is phosphorylated at Thr18 and Ser19 by myosin light chain kinase (MLCK) in a Ca²⁺/calmodulin-dependent manner (1). This phosphorylation is correlated with myosin ATPase activity and smooth muscle contraction (2). ROCK also phosphorylates Ser19 of smooth muscle MLC2, which regulates the assembly of stress fibers (3), Phosphorylation of smooth muscle MLC2 at Ser1/Ser2 and Ser9 by PKC and cdc2 has been reported to inhibit myosin ATPase activity (4,5). Phosphorylation by cdc2 controls the timing of cytokinesis (5). Transgenic mice lacking phosphorylation sites on the cardiac muscle isoform show morphological and

functional abnormalities (6).

Background References

1. Ikebe, M. and Hartshorne, D.J. (1985) J. Biol. Chem. 260, 10027-10031.

2. Tan, J. L. et al. (1992) Annu. Rev. Biochem. 61, 721-759.

3. Totsukawa, G. et al. (2000) J. Cell Biol. 150, 797-806.

4. Ikebe, M. et al. (2000) J. Biol. Chem. 262, 9569-9573.

5. Satterwhite, L. L. et al. (1992) J. Cell Biol. 118, 595-605.

6. Sanbe, A. et al. (1999) J. Biol. Chem. 274, 21085-21094.

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS,

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key WB: Western Blotting

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster **Cross-Reactivity Key**

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Myosin Light Chain 2 Antibody (#3672) Datasheet Without Images Cell Signaling Technology

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.