

#3757 Store at -20°C

SNIP/p140Cap Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M R	Endogenous	145	Rabbit	#Q9C0H9	80725

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

SNIP/p140Cap Antibody detects endogenous levels of total SNIP/p140Cap protein.

Species predicted to react based on 100% sequence homology:

Monkey

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human SNIP/p140Cap. Antibodies were purified by protein A and peptide affinity chromatography.

Background

SNIP (SNAP25-interacting protein)/p140Cap (p130Cas-associated protein) is a cytoskeleton-associated protein identified initially in rat as a protein interacting with the brain-specific synaptosome protein SNAP25 (1) and subsequently as interacting with the broadly expressed scaffold protein p130Cas (2). SNAP25, a presynaptic protein implicated in neurotransmitter secretion, membrane fusion and neurite outgrowth, is part of the SNARE complex that includes syntaxin and synaptobrevin/VAMP (3). SNIP-SNAP25 association is mediated by coiled-coil interactions (1). Overexpression of SNIP inhibits calcium-dependent exocytosis in PC12 cells (1). Human and mouse orthologs of SNIP, termed p140Cap, were subsequently identified through association with p130Cas, a substrate of v-Src and v-Crk that is tyrosine-phosphorylated in response to cell adhesion and mitogenic stimuli (2,4,5). Expression of p140Cap was observed in brain, testis and epithelial-rich tissues and may exist in various alternatively spliced, tissue-specific isoforms (2). p140Cap is also tyrosine-phosphorylated in response to adhesion molecules and EGF treatment (2). Together these studies suggest a role for SNIP/p140Cap in controlling cell spreading, migration and neurosecretion.

Background References

- Chin, L.S. et al. (2000) *J. Biol. Chem.* 275, 1191-1200.
- Di Stefano, P. et al. (2004) *Mol. Biol. Cell* 15, 787-800.
- Gerst, J.E. (1999) *Cell. Mol. Life Sci.* 55, 707-734.
- Sakai, R. et al. (1994) *EMBO J.* 13, 3748-3756.
- Defilippi, P. et al. (2006) *Trends Cell Biol.* 16, 257-263.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Limited Uses

SNIP/p140Cap Antibody (#3757) Datasheet Without Images Cell Signaling Technology

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.