

#3766 Store at -20°C

Thymidylate Synthase Antibody


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA
For Research Use Only. Not for Use in Diagnostic Procedures.

| Applications: | Reactivity: | Sensitivity: | MW (kDa): | Source: | UniProt ID: | Entrez-Gene Id: |
|---------------|-------------|--------------|-----------|---------|-------------|-----------------|
| WB | H M R Hm Mk | Endogenous | 30 | Rabbit | #P04818 | 7298 |

| | | |
|----------------------------------|---|---------------------------|
| Product Usage Information | Application Western Blotting | Dilution 1:1000 |
| Storage | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody. | |
| Specificity / Sensitivity | Thymidylate Synthase Antibody detects endogenous levels of total thymidylate synthase protein. | |
| Source / Purification | Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human thymidylate synthase. Antibodies are purified using protein A and peptide affinity chromatography. | |
| Background | <p>The methylation of deoxyuridine monophosphate (dUMP) to deoxythymidine monophosphate (dTMP) is an essential step in the formation of thymine nucleotides (1,2, reviewed in 3). This process is catalyzed by thymidylate synthase (TS or TYMS), a homodimer composed of two 30 kDa subunits. TS is an intracellular enzyme that provides the sole <i>de novo</i> source of thymidylate, making it a required enzyme in DNA biosynthesis with activity highest in proliferating cells (1). Being the exclusive source of dTMP, investigators have concluded that TS is also an important target for anticancer agents such as 5-fluorouracil (5-FU) (1-5). 5-FU acts as a TS inhibitor and is active against solid tumors such as colon, breast, head, and neck. Research studies have demonstrated that patients with metastases expressing lower levels of TS have a higher response rate to treatment with 5-FU than patients with tumors that have increased levels of TS (5). Researchers continue to investigate TS expression in different types of cancers (6-10).</p> | |
| Background References | <ol style="list-style-type: none"> 1. Johnston, P.G. et al. (1991) <i>Cancer Res</i> 51, 6668-76. 2. Aschele, C. et al. (2002) <i>Ann Oncol</i> 13, 1882-92. 3. Jackman, A.L. and Calvert, A.H. (1995) <i>Ann Oncol</i> 6, 871-81. 4. Van Triest, B. et al. (2000) <i>J Histochem Cytochem</i> 48, 755-60. 5. Johnston, P.G. et al. (1994) <i>J Clin Oncol</i> 12, 2640-7. 6. Kwon, H.C. et al. (2007) <i>Ann Oncol</i> 18, 504-9. 7. Allegra, C.J. et al. (2002) <i>J Clin Oncol</i> 20, 1735-43. 8. Allegra, C.J. et al. (2003) <i>J Clin Oncol</i> 21, 241-50. 9. Tsourouffis, G. et al. (2008) <i>Dig Dis Sci</i> 53, 1289-96. 10. Kim, S.H. et al. (2009) <i>Am J Clin Oncol</i> 32, 38-43. | |

| | |
|-------------------------------|--|
| Species Reactivity | Species reactivity is determined by testing in at least one approved application (e.g., western blot). |
| Western Blot Buffer | IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight. |
| Applications Key | WB: Western Blotting |
| Cross-Reactivity Key | H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected |
| Trademarks and Patents | <p>Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.</p> <p>All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.</p> |
| Limited Uses | |

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.