Plexin A4 (C5D1) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Applications: WB, IP	Reactivity: M R	Sensitivity: Endogenous	MW (kDa): 212	Source/Isotype: Rabbit IgG	UniProt ID: #Q9HCM2	Entrez-Gene Id 91584
Product Usage Information	Ар	plication			Dilution	
	We	stern Blotting			1:1000	
	Imr	munoprecipitation			1:50	
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mN 0.02% sodium azide. Store at -20° C. Do not aliquot to					erol and less than
Specificity / Sensitivity		Plexin A4 (C5D1) Rabbit mAb detects endogenous levels of total Plexin A4 protein.				
Species predicte react based on 10 sequence homolo	00%	nan				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a GST-fusion protein corresponding to residues from the human sequence of Plexin A4.				
Background		Class 3 secreted semaphorin (Sema3A) is a chemorepellent that acts upon a wide variety of axons. As such, it induces a dramatic redistribution and depolymerization of actin filaments that results in growth cone collapse. Plexins are single membrane-spanning signaling proteins encompassing Plexin A1, A2, A3, and A4. Plexins form a complex with neuropilin-1 and -2 and the cell adhesion protein L1 to form a functional semaphorin receptor (1,2). The GTPase Rnd1 binds to the cytoplasmic domain of Plexin A1 to trigger cytoskeletal collapse. In contrast, the GTPase RhoD blocks Rnd1-mediated Plexin A1 activation and repulsion of sympathetic axons by Sema3A (3). Both Plexin A3 and A4 mediate the responses to class 3 Semaphorins in sensory and sympathetic neurons. In particular, Plexin A4 is responsible for signaling of Sema3A via neuropilin-1, while Plexin A3 is				

responsible for signaling of Sema3F via neuropilin-2 (4). 1. Pasterkamp, R.J. and Kolodkin, A.L. (2003) Curr Opin Neurobiol 13, 79-89.

2. Takahashi, T. and Strittmatter, S.M. (2001) Neuron 29, 429-39. 3. Zanata, S.M. et al. (2002) J Neurosci 22, 471-7.

4. Yaron, A. et al. (2005) Neuron 45, 513-23.

Species reactivity is determined by testing in at least one approved application (e.g., western blot). **Species Reactivity**

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, **Western Blot Buffer**

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

WB: Western Blotting IP: Immunoprecipitation **Applications Key**

Cross-Reactivity Key H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Background References

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses

Plexin A4 (C5D1) Rabbit mAb (#3816) Datasheet Without Images Cell Signaling Technology

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.