Store at -200

DDR1 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 125	Source: Rabbit	UniProt ID: #Q08345	Entrez-Gene Id: 780	
Product Usage Information	Application			Dilution			
	We	stern Blotting			1:1000		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity / Sensitiv		DDR1 Antibody detects endogenous levels of total DDR1 protein. The antibody does not cross-react with other proteins of the DDR family.					
Source / Purification	resi	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly598 of human DDR1 protein. Antibodies are purified by protein A and peptide affinity chromatography.					
Background	theii DDF hav abe	The discoidin domain receptors (DDRs) are receptor tyrosine kinases with a discoidin homology repeat in their extracellular domains, activated by binding to extracellular matrix collagens. So far, two mammalian DDRs have been identified: DDR1 and DDR2 (1). They are widely expressed in human tissues and may have roles in smooth muscle cell-mediated collagen remodeling (2). Research studies have implicated aberrant expression and signaling of DDRs in human diseases related to increased matrix degradation and remodeling, such as cardiovascular disease, liver fibrosis, and tumor invasion (1).					
Background Refere	nces 1. V	1. Vogel, W. (1999) <i>FASEB J</i> 13 Suppl, S77-82.					

2. Ferri, N. et al. (2004) Am J Pathol 164, 1575-85.

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry

milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

WB: Western Blotting **Applications Key**

Cross-Reactivity Key H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dq: dog Pq: piq Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. Trademarks and **Patents**

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the **Limited Uses** following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse

1/1/24, 9:12 AM

DDR1 Antibody (#3917) Datasheet Without Images Cell Signaling Technology

engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.