#4021 Store at -20C

## Talin-1 (C45F1) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:Reactivity:Sensitivity:MW (kDa):Source/Isotype:UniProt ID:Entrez-Gene Id:WB, IHC-PH M R Hm MkEndogenous270Rabbit IgG#Q9Y4907094

Product Usage<br/>InformationApplicationDilution<br/>1:1000Western Blotting<br/>Immunohistochemistry (Paraffin)1:50

Storage Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than

0.02% sodium azide. Store at  $-20^{\circ}$ C. Do not aliquot the antibody.

For a carrier free (BSA and azide free) version of this product see product #30960.

Specificity / Sensitivity Talin-1 (C45F1) Rabbit mAb recognizes endogenous levels of total Talin-1 protein.

**Source / Purification**Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to

residues surrounding Ala1772 of human Talin-1 protein.

**Background** Focal adhesions connect the cytoskeleton with the extracellular matrix (ECM), a complex structure of

secreted macromolecules that surrounds mammalian organs and tissues. Integrins clustered on the extracellular side of focal adhesions signal from the ECM to intracellular protein complexes, which in turn signal to the actin cytoskeleton to regulate the tension needed for cell motility. Internal signals also converge on focal adhesions to regulate integrin affinity and avidity. Signaling through focal adhesions regulates cell adhesion, migration, proliferation, apoptosis, and gene expression, and impacts cellular processes such as development, wound healing, immune response, invasion, metastasis, and angiogenesis (reviewed in 1-3). Talin is a large, multidomain focal adhesion protein that interacts with the intracellular domains of integrins and other focal adhesion proteins. Talin is involved in the formation of focal adhesions and in linking focal adhesions to the actin cytoskeleton (4). The interaction between talin

focal adhesions and in linking focal adhesions to the actin cytoskeleton (4). The interaction between talin and integrins increases the affinity between integrin and both insoluble and soluble ECM proteins (5,6).

Background References 1. Burridge, K. et al. (1988) Annu Rev Cell Biol 4, 487-525.

2. Calderwood, D.A. et al. (2000) J Biol Chem 275, 22607-10.

3. ffrench-Constant, C. and Colognato, H. (2004) Trends Cell Biol 14, 678-86.

4. Nayal, A. et al. (2004) Curr Opin Cell Biol 16, 94-8.

5. Tanentzapf, G. and Brown, N.H. (2006) Nat Cell Biol 8, 601-6.

6. Tadokoro, S. et al. (2003) Science 302, 103-6.

**Species Reactivity** Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS,

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key WB: Western Blotting IHC-P: Immunohistochemistry (Paraffin)

Cross-Reactivity Key H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

**Limited Uses** 

Talin-1 (C45F1) Rabbit mAb (#4021) Datasheet Without Images Cell Signaling Technology Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.