

#4043 Store at -20C

# BAP31 Antibody



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R Mk	Endogenous	28	Rabbit	#P51572-2	10134

## Product Usage Information

### Application

Western Blotting  
Immunoprecipitation

### Dilution

1:1000  
1:25

## Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

## Specificity / Sensitivity

BAP31 Antibody detects endogenous levels of total BAP31 protein.

## Species predicted to react based on 100% sequence homology:

Bovine, Dog

## Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to a region surrounding Arg71 of BAP31. Antibodies were purified by peptide affinity chromatography.

## Background

BAP31 (B-cell receptor-association protein 31) is a transmembrane protein associated with the endoplasmic reticulum (ER) and ER-Golgi intermediates and has been implicated in protein trafficking and apoptosis (1,2). During apoptosis Bap31 is cleaved by caspase-8 at two carboxy-terminal sites which can then direct apoptotic signals between the ER and mitochondria (2-4). Association of BAP31 with the anti-apoptotic proteins Bcl-2 or Bcl-xL could function to regulate this ER-mitochondrial pathway (2,5). Several studies have shown that BAP31 can control the trafficking of select proteins between the ER and Golgi apparatus and can affect the transport of proteins to the cell surface (6-10).

## Background References

- Adachi, T. et al. (1996) *EMBO J* 15, 1534-41.
- Ng, F.W. et al. (1997) *J Cell Biol* 139, 327-38.
- Breckenridge, D.G. et al. (2003) *J Cell Biol* 160, 1115-27.
- Chandra, D. et al. (2004) *Mol Cell Biol* 24, 6592-607.
- Ng, F.W. and Shore, G.C. (1998) *J Biol Chem* 273, 3140-3.
- Annaert, W.G. et al. (1997) *J Cell Biol* 139, 1397-410.
- Lambert, G. et al. (2001) *J Biol Chem* 276, 20340-5.
- Schamel, W.W. et al. (2003) *Proc Natl Acad Sci USA* 100, 9861-6.
- Paquet, M.E. et al. (2004) *J Immunol* 172, 7548-55.
- Szczesna-Skorupa, E. and Kemper, B. (2006) *J Biol Chem* 281, 4142-8.

## Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

## Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

## Applications Key

**WB:** Western Blotting **IP:** Immunoprecipitation

## Cross-Reactivity Key

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse **GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](https://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.