

#4165  
Store at -20C

## Pim-3 (D17C9) Rabbit mAb



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H M R	Endogenous	35	Rabbit IgG	#Q86V86	415116

<b>Product Usage Information</b>	<b>Application</b> Western Blotting	<b>Dilution</b> 1:1000
<b>Storage</b>	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
<b>Specificity / Sensitivity</b>	Pim-3 (D17C9) Rabbit mAb detects endogenous levels of total Pim-3 protein. It does not cross-react with other Pim family members.	
<b>Species predicted to react based on 100% sequence homology:</b>	Monkey	
<b>Source / Purification</b>	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ser275 of human Pim-3.	

## Background

### Pim-3 (D17C9) Rabbit mAb (#4165) Datasheet Without Images Cell Signaling Technology

Pim proteins (Pim-1, Pim-2 and Pim-3) are oncogene-encoded serine/threonine kinases (1). Pim-1, a serine/threonine kinase highly expressed in hematopoietic cells, plays a critical role in the transduction of mitogenic signals and is rapidly induced by a variety of growth factors and cytokines (1-4). Pim-1 cooperates with c-Myc in lymphoid cell transformation and protects cells from growth factor withdrawal and genotoxic stress-induced apoptosis (5,6). Pim-1 also enhances the transcriptional activity of c-Myb through direct phosphorylation within the c-Myb DNA binding domain as well as phosphorylation of the transcriptional coactivator p100 (7,8). Hypermutations of the Pim-1 gene are found in B-cell diffuse large cell lymphomas (9). Phosphorylation of Pim-1 at Tyr218 by Etk occurs following IL-6 stimulation and correlates with an increase in Pim-1 activity (10). Various Pim substrates have been identified; Bad is phosphorylated by both Pim-1 and Pim-2 at Ser112 and this phosphorylation reverses Bad-induced cell apoptosis (11,12).

Pim-3, originally named Kid-1, was identified as a protein induced by depolarization in PC-12 cells (13). Like other family members, Pim-3 can phosphorylate Bad to inhibit Bad-mediated apoptosis (14). Aberrant expression of Pim-3 has been observed in a number of types of cancer, including liver, pancreas, colon and gastric cancer (14-17).

## Background References

1. Mikkers, H. et al. (2004) *Mol Cell Biol* 24, 6104-15.
2. Selten, G. et al. (1986) *Cell* 46, 603-11.
3. Meeker, T.C. et al. (1987) *J Cell Biochem* 35, 105-12.
4. Dautry, F. et al. (1988) *J Biol Chem* 263, 17615-20.
5. Möröy, T. et al. (1993) *Proc Natl Acad Sci USA* 90, 10734-8.
6. Lilly, M. and Kraft, A. (1997) *Cancer Res* 57, 5348-55.
7. Levenson, J.D. et al. (1998) *Mol Cell* 2, 417-25.
8. Winn, L.M. et al. (2003) *Cell Cycle* 2, 258-62.
9. Pasqualucci, L. et al. (2001) *Nature* 412, 341-6.
10. Kim, O. et al. (2004) *Oncogene* 23, 1838-44.
11. Aho, T.L. et al. (2004) *FEBS Lett* 571, 43-9.
12. Yan, B. et al. (2003) *J Biol Chem* 278, 45358-67.
13. Feldman, J.D. et al. (1998) *J Biol Chem* 273, 16535-43.
14. Li, Y.Y. et al. (2006) *Cancer Res* 66, 6741-7.
15. Fujii, C. et al. (2005) *Int J Cancer* 114, 209-18.
16. Popivanova, B.K. et al. (2007) *Cancer Sci* 98, 321-8.
17. Zheng, H.C. et al. (2008) *J Cancer Res Clin Oncol* 134, 481-8.

## Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

## Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

## Applications Key

**WB:** Western Blotting

## Cross-Reactivity Key

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse **GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

## Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • [orders@cellsignal.com](mailto:orders@cellsignal.com) • Support: 877-678-TECH (8324) • [info@cellsignal.com](mailto:info@cellsignal.com) • Web: [cellsignal.com](http://cellsignal.com)  
For Research Use Only. Not for Use in Diagnostic Procedures.