

**#4208** Store at -20°C

## Phospho-NMDA Receptor 2B (GluN2B) (Tyr1472) Antibody


**Cell Signaling**  
TECHNOLOGY®

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> WB, IP	<b>Reactivity:</b> R	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 190	<b>Source:</b> Rabbit	<b>UniProt ID:</b> #Q13224	<b>Entrez-Gene Id:</b> 2904
--------------------------------	-------------------------	-----------------------------------	-------------------------	--------------------------	-------------------------------	--------------------------------

<b>Product Usage Information</b>	<b>Application</b> Western Blotting Immunoprecipitation	<b>Dilution</b> 1:1000 1:50
<b>Storage</b>	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
<b>Specificity / Sensitivity</b>	Phospho-NMDA Receptor 2B (GluN2B) (Tyr1472) Antibody detects endogenous levels of NMDA Receptor 2B (GluN2B) only when phosphorylated at Tyr1472.	
<b>Species predicted to react based on 100% sequence homology:</b>	Human, Mouse	
<b>Source / Purification</b>	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr1472 of NMDA Receptor 2B (GluN2B). Antibodies are purified by protein A and peptide affinity chromatography.	
<b>Background</b>	<p>N-methyl-D-aspartate receptor (NMDAR) forms a heterodimer of at least one NR1 and one NR2A-D subunit. Multiple receptor isoforms with distinct brain distributions and functional properties arise by selective splicing of the NR1 transcripts and differential expression of the NR2 subunits. NR1 subunits bind the co-agonist glycine and NR2 subunits bind the neurotransmitter glutamate. Activation of the NMDA receptor or opening of the ion channel allows flow of Na<sup>+</sup> and Ca<sup>2+</sup> ions into the cell, and K<sup>+</sup> out of the cell (1). Each subunit has a cytoplasmic domain that can be directly modified by the protein kinase/phosphatase (2). PKC can phosphorylate the NR1 subunit (NMDAR1) of the receptor at Ser890/Ser896, and PKA can phosphorylate NR1 at Ser897 (3). The phosphorylation of NR1 by PKC decreases its affinity for calmodulin, thus preventing the inhibitory effect of calmodulin on NMDAR (4). The phosphorylation of NR1 by PKA probably counteracts the inhibitory effect of calcineurin on the receptor (5). NMDAR mediates long-term potentiation and slow postsynaptic excitation, which play central roles in learning, neurodevelopment, and neuroplasticity (6).</p> <p>EphrinB2 binding to the receptor EphB leads to the activation of Src family tyrosine kinases, which phosphorylate NMDAR2B at Tyr1252, Tyr1336 and Tyr1472. In turn, phosphorylated NMDAR2B enhances the ability of the functional NMDA receptor to regulate Ca<sup>2+</sup> influx in response to glutamate (7, 8). Phosphorylation of NMDAR2B at Tyr1472 was independently identified at Cell Signaling Technology (CST) using PhosphoScan®, CST's MS/MS platform for phosphorylation site discovery. Phosphorylation of NMDAR2B at Tyr1472 was observed in extracts isolated from ischemic rat brain. For additional information please visit PhosphoSitePlus®, CST's modification site knowledgebase, at <a href="http://www.phosphosite.org">www.phosphosite.org</a>.</p>	
<b>Background References</b>	<ol style="list-style-type: none"> <li>Liu, X.B. et al. (2004) <i>J Neurosci</i> 24, 8885-95.</li> <li>Westphal, R.S. et al. (1999) <i>Science</i> 285, 93-6.</li> <li>Tingley, W.G. et al. (1997) <i>J Biol Chem</i> 272, 5157-66.</li> <li>Hisatsune, C. et al. (1997) <i>J Biol Chem</i> 272, 20805-10.</li> <li>Raman, I.M. et al. (1996) <i>Neuron</i> 16, 415-21.</li> <li>Makhinson, M. et al. (1999) <i>J Neurosci</i> 19, 2500-10.</li> <li>Nakazawa, T. et al. (2001) <i>J Biol Chem</i> 276, 693-9.</li> <li>Takasu, M.A. et al. (2002) <i>Science</i> 295, 491-5.</li> </ol>	

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**WB:** Western Blotting **IP:** Immunoprecipitation

**Cross-Reactivity Key**

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.