Store at -200

## nNOS (C7D7) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

| <b>Applications:</b><br>WB, IP, IHC-P, IF-F | Reactivity:<br>H M R                | Sensitivity:<br>Endogenous   | <b>MW (kDa):</b><br>160 | Source/Isotype:<br>Rabbit IgG | UniProt ID:<br>#P29475 | Entrez-Gene Id<br>4842 |  |
|---|-------------------------------------|--|-------------------------|-------------------------------|------------------------|------------------------|--|
| Product Usage<br>Information                | Ap                                  | Application  |                         |                               | Dilution               |                        |  |
|   | We                                  | Western Blotting   |                         |                               |                        | 1:1000                 |  |
|   | Imi                                 | Immunoprecipitation  |                         |                               |                        | 1:50                   |  |
|   | Imi                                 | munohistochemistry   | (Paraffin)              | 1:200                         |                        |                        |  |
|   | Imi                                 | munofluorescence (   | Frozen)                 | 1:200                         |                        |                        |  |
| Storage                                     | •                                   | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at $-20^{\circ}$ C. Do not aliquot the antibody.  |                         |                               |                        |                        |  |
|   | For                                 | For a carrier free (BSA and azide free) version of this product see product #67860.  |                         |                               |                        |                        |  |
| Specificity / Sensitiv                      | vity nNC                            | nNOS (C7D7) Rabbit mAb detects endogenous levels of total nNOS protein.  |                         |                               |                        |                        |  |
| Source / Purification                       |                                     | Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ser280 of human nNOS.   |                         |                               |                        |                        |  |
| Background                                  | oxy<br>four<br>lipo<br>exp<br>incli | Nitric Oxide Synthase (NOS) catalyzes the formation of nitric oxide (NO) and citrulline from L-arginine, oxygen, and cofactors. Three family members have been characterized: neuronal NOS (nNOS), which is found primarily in neuronal tissue; inducible NOS (iNOS), which is induced by interferon gamma and lipopolysaccharides in the kidney and cardiovascular system; and endothelial NOS (eNOS), which is expressed in blood vessels (1). NO is a messenger molecule with diverse functions throughout the body, including the maintenance of vascular integrity, homeostasis, synaptic plasticity, long-term potentiation, learning, and memory (2,3). |                         |                               |                        |                        |  |
| Background Refere                           | 2. S                                | <ol> <li>Tsutsui, M. (2004) J Atheroscler Thromb 11, 41-8.</li> <li>Son, H. et al. (1996) Cell 87, 1015-23.</li> <li>Hawkins, R.D. (1996) Neuron 16, 465-7.</li> </ol>   |                         |                               |                        |                        |  |

Species reactivity is determined by testing in at least one approved application (e.g., western blot). **Species Reactivity** 

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, Western Blot Buffer

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

WB: Western Blotting IP: Immunoprecipitation IHC-P: Immunohistochemistry (Paraffin) **Applications Key** 

IF-F: Immunofluorescence (Frozen)

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster **Cross-Reactivity Key** 

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Cell Signaling Technology is a trademark of Cell Signaling Technology. Inc. Trademarks and **Patents** 

U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the **Limited Uses** 

following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

nNOS (C7D7) Rabbit mAb (#4231) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.