

#4337 Store at -20C

GSK-3 α (D80E6) Rabbit mAb


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R Hm Mk	Endogenous	51	Rabbit IgG	#P49840	2931

Product Usage Information	Application Western Blotting Immunoprecipitation	Dilution 1:1000 1:50
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C . Do not aliquot the antibody.	
Specificity / Sensitivity	GSK-3 α (D80E6) Rabbit mAb detects endogenous levels of total GSK-3 α protein. The antibody does not cross-react with GSK-3 β .	
Species predicted to react based on 100% sequence homology:	Pig	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to the sequence of human GSK-3 α protein.	
Background	Glycogen synthase kinase-3 (GSK-3) was initially identified as an enzyme that regulates glycogen synthesis in response to insulin (1). GSK-3 is a ubiquitously expressed serine/threonine protein kinase that phosphorylates and inactivates glycogen synthase. GSK-3 is a critical downstream element of the PI3K/Akt cell survival pathway whose activity can be inhibited by Akt-mediated phosphorylation at Ser21 of GSK-3 α and Ser9 of GSK-3 β (2,3). GSK-3 has been implicated in the regulation of cell fate in <i>Dictyostelium</i> and is a component of the Wnt signaling pathway required for <i>Drosophila</i> , <i>Xenopus</i> , and mammalian development (4). GSK-3 has been shown to regulate cyclin D1 proteolysis and subcellular localization (5). GSK-3 α regulates the production of amyloid- β peptides, a major component of the plaques that accumulate with progression of Alzheimer disease. Administration of therapeutic concentrations of lithium, a GSK-3 inhibitor, attenuates amyloid- β production by specifically inhibiting the cleavage of amyloid precursor protein (APP) by γ -secretase, blocking accumulation of amyloid- β peptides in the brains of mice that overproduce APP (6).	
Background References	1. Welsh, G.I. et al. (1996) <i>Trends Cell Biol</i> 6, 274-9. 2. Srivastava, A.K. and Pandey, S.K. (1998) <i>Mol Cell Biochem</i> 182, 135-41. 3. Cross, D.A. et al. (1995) <i>Nature</i> 378, 785-9. 4. Nusse, R. (1997) <i>Cell</i> 89, 321-3. 5. Diehl, J.A. et al. (1998) <i>Genes Dev</i> 12, 3499-511. 6. Phiel, C.J. et al. (2003) <i>Nature</i> 423, 435-439.	
Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.	
Applications Key	WB: Western Blotting IP: Immunoprecipitation	
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected	

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.