Store at -20C

p44 MAP Kinase (Erk1) Antibody



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support:

877-678-TECH (8324)

Web:

info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Applications: WB, IP	Reactivity: M R	Sensitivity: Endogenous	MW (kDa): 44	Source: Rabbit	UniProt ID: #P27361	Entrez-Gene Id: 5595	
Product Usage Information	Ap	Application			Dilution		
	We	Western Blotting			1:1000		
	Imi	Immunoprecipitation			1:50		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity / Sensiti		p44 MAP Kinase (Erk1) Antibody detects endogenous levels of p44 MAP kinase (Erk1). This antibody does not recognize p38 MAP kinase, p42 MAP kinase (Erk2) or SAPK/JNK.					
Source / Purificatio	N-te	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the N-terminus of p44 MAP kinase (Erk1). Antibodies are purified by protein A and peptide affinity chromatography.					
Background	kina The extr con thre MAI	Mitogen-activated protein kinases (MAPKs) are a widely conserved family of serine/threonine protein kinases involved in many cellular programs, such as cell proliferation, differentiation, motility, and death. The p44/42 MAPK (Erk1/2) signaling pathway can be activated in response to a diverse range of extracellular stimuli, including mitogens, growth factors, and cytokines (1-3), and research investigators consider it an important target in the diagnosis and treatment of cancer (4). Upon stimulation, a sequential three-part protein kinase cascade is initiated, consisting of a MAP kinase kinase (MAPKKK or MAP3K), a MAP kinase kinase (MAPKK or MAP2K), and a MAP kinase (MAPK). Multiple p44/42 MAP3Ks have been identified, including members of the Raf family, as well as Mos and Tpl2/COT. MEK1 and MEK2					

Background References

1. Roux, P.P. and Blenis, J. (2004) Microbiol Mol Biol Rev 68, 320-44.

as DUSPs or MKPs (10), along with MEK inhibitors, such as U0126 and PD98059.

- 2. Baccarini, M. (2005) FEBS Lett 579, 3271-7.
- 3. Meloche, S. and Pouysségur, J. (2007) Oncogene 26, 3227-39.
- 4. Roberts, P.J. and Der, C.J. (2007) Oncogene 26, 3291-310.
- 5. Rubinfeld, H. and Seger, R. (2005) Mol Biotechnol 31, 151-74.
- 6. Murphy, L.O. and Blenis, J. (2006) Trends Biochem Sci 31, 268-75.
- 7. Dalby, K.N. et al. (1998) J Biol Chem 273, 1496-505.
- 8. Marais, R. et al. (1993) Cell 73, 381-93.
- 9. Kortenjann, M. et al. (1994) Mol Cell Biol 14, 4815-24.
- 10. Owens, D.M. and Keyse, S.M. (2007) Oncogene 26, 3203-13.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

are the primary MAPKKs in this pathway (5,6). MEK1 and MEK2 activate p44 and p42 through phosphorylation of activation loop residues Thr202/Tyr204 and Thr185/Tyr187, respectively. Several downstream targets of p44/42 have been identified, including p90RSK (7) and the transcription factor Elk-1 (8,9), p44/42 are negatively regulated by a family of dual-specificity (Thr/Tyr) MAPK phosphatases, known

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

3/23/24. 1:14 PM

Trademarks and Patents

Limited Uses

p44 MAP Kinase (Erk1) Antibody (#4372) Datasheet Without Images Cell Signaling Technology

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.