

#4636 Store at -20C

MAD2L1 (D8A7) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H M R	Endogenous	22	Rabbit IgG	#Q13257	4085

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

MAD2L1 (D8A7) Rabbit mAb detects endogenous levels of total MAD2L1 protein.

Species predicted to react based on 100% sequence homology:

Monkey, Bovine, Dog

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues at the amino terminus of human MAD2L1 protein.

Background

Correct segregation of sister chromatids prior to the onset of cell division is essential to the maintenance of genetic integrity and the avoidance of aneuploidy and chromosomal instability, characteristics of many cancer cells. The mitotic checkpoint, also known as the spindle assembly checkpoint, monitors accurate attachment of kinetochores to the spindle, inhibits mitosis and delays the onset of anaphase until all chromosomes are aligned at the metaphase plate (1). MAD2L1 is an essential participant in the mitotic checkpoint (2). It exists in two conformations, including the open and inactive O-MAD2 form and the closed, active C-MAD2 form. Prior to mitosis, MAD2L1 is localized to the cytosol and exists largely in the closed, inactive form. During the mitotic checkpoint, MAD2L1 switches to the open, active conformation (3). Together with other checkpoint proteins, MAD2L1 binds to and deactivates Cdc20, thereby inhibiting the anaphase promoting complex (4). When the kinetochores are correctly attached to the spindle, MAD2L1 releases Cdc20, which allows activation of the anaphase promoting complex and subsequent degradation of key mitotic substrates and the initiation of metaphase-anaphase transition (5).

Background References

1. Wassmann, K. and Benezra, R. (2001) *Curr Opin Genet Dev* 11, 83-90.
2. Musacchio, A. and Salmon, E.D. (2007) *Nat Rev Mol Cell Biol* 8, 379-93.
3. Skinner, J.J. et al. (2008) *J Cell Biol* 183, 761-8.
4. Reddy, S.K. et al. (2007) *Nature* 446, 921-5.
5. Stegmeier, F. et al. (2007) *Nature* 446, 876-81.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse **GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses**MAD2L1 (D8A7) Rabbit mAb (#4636) Datasheet Without Images Cell Signaling Technology**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.