

#4759 Store at -20°C

ER Protein Folding Antibody Sampler Kit

1 Kit (7 x 20 microliters)



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
BiP (C50B12) Rabbit mAb	3177	20 µl	78 kDa	Rabbit IgG
Ero1-L α Antibody	3264	20 µl	60 kDa	Rabbit
ERp44 (D17A6) XP® Rabbit mAb	3798	20 µl	44 kDa	Rabbit IgG
ERp57 (G117) Antibody	2881	20 µl	57 kDa	Rabbit
ERp72 (D70D12) XP® Rabbit mAb	5033	20 µl	72 kDa	Rabbit IgG
Grp94 Antibody	2104	20 µl	100 kDa	Rabbit
PDI (C81H6) Rabbit mAb	3501	20 µl	57 kDa	Rabbit
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

The ER Protein Folding Antibody Sampler Kit contains reagents to investigate the initiation of translation within the cell. The kit contains enough primary and secondary antibodies to perform two Western blot experiments per primary antibody.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Background

After their synthesis, secretory proteins translocate into the endoplasmic reticulum (ER) where they are post-translationally modified and properly folded. To reach their native conformation, many secretory proteins require the formation of intra- or inter-molecular disulfide bonds (1). This process is called oxidative protein folding. Disulfide isomerase (PDI) catalyzes the formation and isomerization of these disulfide bonds (2). Studies on mechanisms of oxidative folding suggest that molecular oxygen oxidizes the ER-protein Ero1, which in turn oxidizes PDI through disulfide exchange (3). This event is then followed by PDI-catalyzed disulfide bond formation on folding proteins (3). Other ER resident proteins that possess the thioredoxin homology domains, including endoplasmic reticulum stress proteins 72, 57 and 44 (ERp72, ERp57 and ERp44), constitute the PDI family (4,5,6). The ER also contains a pool of molecular chaperones, including Grp94, to help proteins fold properly. Grp94 is a glucose-regulated protein (7) with sequence homology to Hsp90 (8). BiP is another chaperone whose synthesis is increased when protein folding is disturbed. BiP binds to misfolded proteins to prevent them from forming aggregates and assists in proper refolding (9).

Background References

1. Huppa, J.B. and Ploegh, H.L. (1998) *Cell* 92, 145-8.
2. Ellgaard, L. and Ruddock, L.W. (2005) *EMBO Rep* 6, 28-32.
3. Tu, B.P. and Weissman, J.S. (2004) *J Cell Biol* 164, 341-6.
4. Mazzarella, R.A. et al. (1990) *J Biol Chem* 265, 1094-101.
5. Satoh, M. et al. (2005) *Cell Stress Chaperones* 10, 278-84.
6. Jessop, C.E. et al. (2007) *EMBO J* 26, 28-40.
7. Lee, A.S. et al. (1981) *Proc Natl Acad Sci U S A* 78, 4922-5.
8. Sorger, P.K. and Pelham, H.R. (1987) *J Mol Biol* 194, 341-4.
9. Kohno, K. et al. (1993) *Mol Cell Biol* 13, 877-90.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.