

#4852 Store at -20°C

PCTAIRE 1 Antibody


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 65	Source: Rabbit	UniProt ID: #Q00536	Entrez-Gene Id: 5127
--------------------------------	-----------------------------	-----------------------------------	------------------------	--------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

PCTAIRE 1 Antibody detects endogenous levels of total PCTAIRE 1.

Species predicted to react based on 100% sequence homology:

Human

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Arg98 of human PCTAIRE 1 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Three distinct PCTAIRE isoforms (PCTAIRE 1, PCTAIRE 2 and PCTAIRE 3) have been identified in humans and belong to the CDK family of serine/threonine protein kinases. These proteins have a core kinase domain flanked by unique amino- and carboxy-terminal domains. CDK proteins are known to regulate the cell cycle. All three PCTAIRE isoforms are abundantly expressed and catalytically active in post-mitotic brain, suggesting that they may function in processes other than cell division (1). PCTAIRE 1 is a cytoplasmic phosphoprotein whose kinase activity peaks in G2 and S phase (2). While one study indicates that noncovalent interactions with a regulatory component (such as a cyclin) are necessary for catalytic activity of PCTAIRE 1, others show that the monomeric protein is fully active (3). The Cdk5/p25 complex phosphorylates PCTAIRE 1 at Ser95, enhancing its kinase activity (4).

Background References

1. Meyerson, M. et al. (1992) *EMBO J.* 11, 2909-2917.
2. Charrasse, S. et al. (1999) *Cell Growth Differ.* 10, 611-620.
3. Graeser, R. et al. (2002) *J. Cell Sci.* 115, 3479-3490.
4. Cheng, K. et al. (2002) *J. Biol. Chem.* 277, 31988-31993.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.