


#4932 Store at -20C

FAF1 Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M R	Endogenous	74 to 80	Rabbit	#Q9UNN5	11124

Product Usage Information	Application Western Blotting	Dilution 1:1000
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	FAF1 Antibody detects endogenous levels of FAF1 protein.	
Source / Purification	Polyclonal antibodies are produced by immunizing animals with synthetic peptides corresponding to a central region of FAF1 around residue 260. Antibodies are purified by protein A and affinity chromatography.	
Background	FAF1 was originally identified through yeast two-hybrid screening, interacting with the cytoplasmic domain of Fas, a member of the TNF receptor superfamily that plays a critical role in in apoptosis during development and immune function (1). FAF1 is widely expressed with highest expression observed in testis, skeletal muscle and heart (2). FAF1 potentiates Fas-mediated apoptosis and may induce apoptosis without Fas stimulation in some cell types. It does not contain typical death motifs, but rather has two amino-terminal domains with structural homology to ubiquitin. While the precise role of FAF1 during apoptosis is still unclear, it has been observed to be one of the components of the death-inducing signaling complex (DISC) during Fas-mediated apoptosis and can bind to caspase-8 and FADD (3). FAF1 has also been shown to suppress the activation of the NF-kappaB transcription factor (4).	
Background References	1. Chu, K. et al. (1995) <i>Proc Natl Acad Sci U S A</i> 92, 11894-8. 2. Ryu, S.W. et al. (1999) <i>Biochem Biophys Res Commun</i> 262, 388-94. 3. Ryu, S.W. et al. (2003) <i>J Biol Chem</i> 278, 24003-10. 4. Park, M.Y. et al. (2004) <i>J Biol Chem</i> 279, 2544-9.	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the</p>

end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.