## **NIK Antibody**



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

#### For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H M	Sensitivity: Endogenous	<b>MW (kDa):</b> 125	Source: Rabbit	UniProt ID: #Q99558	Entrez-Gene Id: 9020	
Product Usage Information	Application			Dilution			
	Western Blotting			1:1000			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity / Sensit	ivity NIK	NIK Antibody detects endogenous levels of total NIK protein.					

Species predicted to react based on 100% sequence homology:

Rat, Monkey, Bovine, Horse

#### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues adjacent to glycine 659 of human NIK. Antibodies are purified by protein A and peptide affinity chromatography.

#### **Background**

Transcription factors of the nuclear factor  $\kappa B$  (NF- $\kappa B$ )/Rel family play a pivotal role in inflammatory and immune responses (1,2). There are five family members in mammals: RelA, c-Rel, RelB, NF- $\kappa B1$  (p105/p50), and NF- $\kappa B2$  (p100/p52). Both p105 and p100 are proteolytically processed by the proteasome to produce p50 and p52, respectively. Rel proteins bind p50 and p52 to form dimeric complexes that bind DNA and regulate transcription. In unstimulated cells, NF- $\kappa B$  is sequestered in the cytoplasm by I $\kappa B$  inhibitory proteins (3-5). NF- $\kappa B$ -activating agents can induce the phosphorylation of I $\kappa B$  proteins, targeting them for rapid degradation through the ubiquitin-proteasome pathway and releasing NF- $\kappa B$  to enter the nucleus where it regulates gene expression (6-8). NIK and IKK $\alpha$  (IKK1) regulate the phosphorylation and processing of NF- $\kappa B2$  (p100) to produce p52, which translocates to the nucleus (9-11). Activation of NF-kappaB can be controlled by NF- $\kappa B$ -inducing kinase (NIK), a member of the MAP3K

Activation of NF-kappaB can be controlled by NF-kB-inducing kinase (NIK), a member of the MAP3K family that was originally identified as a TRAF2-interacting protein and thereby coupled to receptor activation (12). NIK forms a complex with and phosphorylates IKK1 and IKK2, subsequently leading to the phosphorylation of IkappaB and translocation of NF-kappaB to the nucleus (13-15).

### **Background References**

- 1. Baeuerle, P.A. and Henkel, T. (1994) Annu Rev Immunol 12, 141-79.
- 2. Baeuerle, P.A. and Baltimore, D. (1996) Cell 87, 13-20.
- 3. Haskill, S. et al. (1991) Cell 65, 1281-9.
- 4. Thompson, J.E. et al. (1995) Cell 80, 573-82.
- 5. Whiteside, S.T. et al. (1997)  $\it EMBO J 16$ , 1413-26.
- 6. Traenckner, E.B. et al. (1995) EMBO J 14, 2876-83.
- 7. Scherer, D.C. et al. (1995) Proc Natl Acad Sci USA 92, 11259-63.
- 8. Chen, Z.J. et al. (1996) Cell 84, 853-62.
- 9. Senftleben, U. et al. (2001) Science 293, 1495-9.
- 10. Coope, H.J. et al. (2002) EMBO J 21, 5375-85.
- 11. Xiao, G. et al. (2001) Mol Cell 7, 401-9.
- 12. Malinin, N.L. et al. (1997) Nature 385, 540-4.
- 13. Régnier, C.H. et al. (1997) Cell 90, 373-83.
- 14. Woronicz, J.D. et al. (1997) Science 278, 866-9.

# 15. Ling, L. et al. (1998) Proc Natl Acad Sci U S A 95, 3792-7.

#### **Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

3/23/24. 11:15 AM

Western Blot Buffer

NIK Antibody (#4994) Datasheet Without Images Cell Signaling Technology

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key** 

**Cross-Reactivity Key** 

WB: Western Blotting

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

**Limited Uses** 

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.