

#5071 Store at -20C

ELP1/IKBKAP Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H Mk	Endogenous	150	Rabbit	#O95163	8518

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

ELP1/IKBKAP Antibody recognizes endogenous levels of total ELP1/IKBKAP protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human ELP1/IKBKAP protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Elongator is a highly conserved transcription elongation factor complex that was first identified in yeast as part of the hyperphosphorylated RNA polymerase II (RNAPII) holoenzyme (1). The Elongator complex consists of 6 subunits, ELP1-6, and has been shown to have acetyltransferase activity (2). The acetylation targets of Elongator include histone H3, which is linked to the transcription elongation function of the complex, and α -tubulin, which is associated with regulation of migration and maturation of projection neurons (3-6).

The ELP1/IKBKAP subunit of Elongator was initially thought to function as a scaffolding protein within the NF κ B signaling pathway (7). It contains several WD40 domains and is critical for the formation of the Elongator complex (8). Investigators have determined that mutations in ELP1 are the cause of Familial Dysautonomia (FD), an autosomal recessive neurodegenerative disorder (9). Research studies have demonstrated that defects in Elongator function upon ELP1 mutation affect transcription elongation of several genes involved in cell motility and neuronal development that may be the underlying cause of the neuropathology of FD patients (10,11).

Background References

- Otero, G. et al. (1999) *Mol Cell* 3, 109-18.
- Creppe, C. and Buschbeck, M. (2011) *J Biomed Biotechnol* 2011, 924898.
- Wittschieben, B.O. et al. (1999) *Mol Cell* 4, 123-8.
- Hawkes, N.A. et al. (2002) *J Biol Chem* 277, 3047-52.
- Kim, J.H. et al. (2002) *Proc Natl Acad Sci USA* 99, 1241-6.
- Creppe, C. et al. (2009) *Cell* 136, 551-64.
- Cohen, L. et al. (1998) *Nature* 395, 292-6.
- Frohloff, F. et al. (2003) *J Biol Chem* 278, 956-61.
- Anderson, S.L. et al. (2001) *Am J Hum Genet* 68, 753-8.
- Close, P. et al. (2006) *Mol Cell* 22, 521-31.
- Cohen-Kupiec, R. et al. (2011) *PLoS One* 6, e19147.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.