#5184 Store at -20C

Phospho-IRF-7 (Ser471/472) Antibody



Orders: 8

877-616-CELL (2355) orders@cellsignal.com

Support: 877-678-TECH (8324)

Web:

info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H	Sensitivity: Transfected Only	MW (kDa): 65	Source: Rabbit	UniProt ID: #Q92985	Entrez-Gene Id 3665
Product Usage Information	Application			Dilution		
	We	Western Blotting			1:1000	
Storage	•	plied in 10 mM sodi C. Do not aliquot the		, 150 mM NaCl, 100 $\mu g/\text{ml}$ BSA and 50% glycerol. Store at $-$		
Specificity / Sensitivity Phospho-IRF-7 (Ser471/472) Antibody and 472.			./472) Antibody det	etects transfected levels of IRF-7 when phosphorylated at Ser471		
Source / Purification	to re	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser471/472 of human IRF-7 protein. Antibodies were purified by protein A and peptide affinity chromatography.				

Background

Interferon regulatory factors (IRFs) comprise a family of transcription factors that function within the Jak/Stat pathway to regulate interferon (IFN) and IFN-inducible gene expression in response to viral infection (1). IRFs play an important role in pathogen defense, autoimmunity, lymphocyte development, cell growth, and susceptibility to transformation. The IRF family includes nine members: IRF-1, IRF-2, IRF-9/ISGF3y, IRF-3, IRF-4 (Pip/LSIRF/ICSAT), IRF-5, IRF-6, IRF-7, and IRF-8/ICSBP. All IRF proteins share homology in their amino-terminal DNA-binding domains. IRF family members regulate transcription through interactions with proteins that share similar DNA-binding motifs, such as IFN-stimulated response elements (ISRE), IFN consensus sequences (ICS), and IFN regulatory elements (IRF-E) (2).

IRF-7, which is functionally similar to IRF-3, is preferentially expressed in lymphoid cells and induced by virus, LPS, and IFN- α (3-5). IRF-7 plays an essential role in the induction of type I interferon in response viral infection (6-8). Like IRF-3, IRF-7 is regulated at multiple serine phosphorylation sites near its carboxyl terminus, which are required for nuclear translocation, DNA binding, and transcriptional activity (9-11).

Background References

- 1. Taniguchi, T. et al. (2001) Annu Rev Immunol 19, 623-55.
- 2. Honda, K. and Taniguchi, T. (2006) Nat Rev Immunol 6, 644-58.
- 3. Au, W.C. et al. (1998) J Biol Chem 273, 29210-7.
- 4. Wathelet, M.G. et al. (1998) *Mol Cell* 1, 507-18.
- 5. Marié, I. et al. (1998) *EMBO J* 17, 6660-9.
- 6. Sato, M. et al. (2000) Immunity 13, 539-48.
- 7. Honda, K. et al. (2005) Nature 434, 772-7.
- 8. Colina, R. et al. (2008) Nature 452, 323-8.
- 9. Lin, R. et al. (2000) J Biol Chem 275, 34320-7.
- 10. Yang, H. et al. (2003) J Biol Chem 278, 15495-504.
- 11. Caillaud, A. et al. (2005) J Biol Chem 280, 17671-7.

Species Reactivity Spe

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

3/23/24. 11:16 AM

Trademarks and Patents

Limited Uses

Phospho-IRF-7 (Ser471/472) Antibody (#5184) Datasheet Without Images Cell Signaling Technology

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.